

Judul Artikel : The Effect of *Strobilanthes crispus* on Blood Glucose Levels and Lipid Profile of Streptozotocin-induced Diabetic Rats
Nama Jurnal : Open Access Macedonian Journal of Medical Sciences
ISSN : 1857-9655
Tahun Terbit : 2022
Penerbit : Scientific Foundation SPIROSKI
Volume : 10
Nomor : T8
No. Halaman : 35-40
Url Cek Plagiasi : http://repositor.almaata.ac.id/id/eprint/3504/1/P5_turnitin.pdf
Url Index : <https://www.scopus.com/sourceid/21100824403>
Url Jurnal : <https://doi.org/10.3889/oamjms.2022.9468>

Rekapitulasi Proses Publikasi

No	Kegiatan	Tanggal
1	Bukti Submisi	21 Maret 2022
2	Pemberitahuan hasil review	21 Maret 2022
3	Pemberitahuan produksi	30 Maret 2022
4	Proofread	20 Juli 2023

Bukti Submisi: 21 Maret 2022



verianiaprilia verianiaprilia <verianiaprilia@almaata.ac.id>

[OAMJMS] Submission Acknowledgement

Prof. Dr Mirko Spiroski via SFS - Journals (Scientific Foundation SPIROSKI - Journals), Skopje, Republic of Macedonia <noreply@publicknowledgeproject.org>

21 Maret 2022
pukul 20.07

Balas Ke: "Prof. Dr Mirko Spiroski" <mspiroski@yahoo.com>

Kepada: Tri Hernawati <Trihernawati87@gmail.com>, Desta Sari D <destasari30@gmail.com>, Indra Kusuma W <26@gmail.com>, Kunti Septiani <kuntiseptiani14@gmail.com>, Hamam Hadi <hhadi@almaata.ac.id>, Yulinda Kurniasari <kurniasari.yulinda@almaata.ac.id>, Effatul Afifah <effatulafifah@almaata.ac.id>, Veriani Aprilia <verianiaprilia@almaata.ac.id>

Hello,

Laras Ayu Fauziah Nur Fitri has submitted the manuscript, "The effect of Strobilanthes crispus on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats " to Open Access Macedonian Journal of Medical Sciences.

[Kutipan teks disembunyikan]

The effect of *Strobilanthes crispus* on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats

Laras Ayu Fauziah Nur Fitri¹, Tri Hernawati¹, Desta Sari D¹, Indra Kusuma W¹, Kunti S¹, Hamam Hadi², Yulinda Kurniasari¹, Effatul Afifah¹, Veriani Aprilia¹

¹Department of Nutrition, Faculty of Health Sciences, Alma Ata University, Jl. Brawijaya 99 Tamantirto, Kasihan, Bantul, Yogyakarta 55183

²Graduate School of Public Health, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta 55183, Indonesia

Corresponding author: Veriani Aprilia, Department of Nutrition, Faculty of Health Sciences, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta 55183, Indonesia. E-mail: verianiaprilia@almaata.ac.id. ORCID iD: 0000-0002-6231-6487

Short title: *Strobilanthes crispus* effect on blood glucose levels and lipid profile

Funding: This research was supported by Universitas Alma Ata

The effect of *Strobilanthes crispus* on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats

Abstract

BACKGROUND: Diabetes mellitus is a metabolic disease associated with unhealthy eating patterns, a lack of physical activity, obesity, smoking habits, and family history. *Strobilanthes crispus* (SC) contains antioxidant compounds known to exert a hypoglycemic effect during long-term interventions in non-diabetic samples.

AIMS: This study sought to determine the effect of administering the extract of SC leaves on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats.

METHODS: A total of 30 male *Rattus norvegicus* rats were divided into control and treatment groups. The control groups consisted of negative (normal rats) and positive controls (diabetic rats), whereas treatment groups consisted of diabetic rats administered with 3.2% and 16.8% SC leaf extract, and glibenclamide as the drug comparison groups. Blood samples were collected from the rats' retroorbital vein for the purposes of glucose level and lipid profile measurement before and after the induction of diabetes with STZ as well as 14 days after the intervention. The collected data were analyzed statistically by means of a one-way analysis of variance (ANOVA) continued with Duncan's multiple range test.

RESULTS: The results showed the rats to be hyperglycemic experienced changes in their lipid profiles following the induction of diabetes. The administration of 3.2% SC leaf extract for 14 days reduced the rats' blood glucose levels, while the effect of the 16.8% SC leaf extract was more pronounced in terms of reducing the rats' blood glucose levels and improving their lipid profiles (reducing the triglyceride, total cholesterol, and low-density lipoprotein cholesterol levels, while increasing the high-density lipoprotein cholesterol levels).

CONCLUSION: This study found 16.8% SC leaf extract to exert a good hypoglycemic effect and lead to an improvement in the lipid profiles in diabetic rats.

Keywords: *Diabetes mellitus, Strobilanthes crispus, blood sugar, lipid profile, rat*

Introduction

Diabetes mellitus (DM) comprises a group of metabolic diseases associated with defects in insulin secretion, insulin action, or a combination of the two (1). DM is currently the fourth leading cause of death worldwide (2). Among the different types of DM, type 2 DM is known to result from unhealthy eating habits, a lack of physical activity, obesity, smoking habits, and/or family history (3).

Hyperglycemia in patients with type 2 DM has been found to increase the formation of free radicals such as the superoxide, hydrogen peroxide, nitric oxide, and hydroxyl

radicals (4). Oxidative stress can result if the levels of free radicals exceed the antioxidants levels in the body, resulting in damage to the cell organelles and enzymes as well as increased lipid peroxidation. Increased levels of fat in the blood (dyslipidemia) serve as an indicator of the occurrence of lipid peroxidation (5). In addition, oxidative stress can also cause complications such as eye disorders, kidney damage, and nerve damage (6).

The consumption of antioxidants could inhibit the activity of free radicals. Keji beling (*Strobilanthes crispus* [SC]) is a local Indonesian plant species known for its bioactive properties, especially when it comes to treating diabetes and improving lipid profiles (i.e., polyphenols, alkaloids, coumarins, flavonoids, iridoids, triterpenes, and sterols) (7)(8). Several in vivo studies involving various doses and formulas of SC have demonstrated these beneficial effects. For example, a prior study found the ethanol extract of SC (a dose of 14.7 mg/g body weight [BW]) to be effective in terms of reducing the blood glucose levels in rats, although the results did not reveal a curative effect in DM subjects [9]. Other studies have examined the effect SC leaf juice administered at doses of 1 mL/kg BW, 1.5 mL/kg BW, and 2 mL/kg BW over a period of 30 days (9) (10) and in the form of ethanol extract administered at doses of 100 mg/kg BW, 200 mg/kg BW, and 400 mg/kg BW over a period of 21 days (11). With regard to the safety of SC leaves, a toxicity analysis was only performed for 14 days of administration, although it revealed no instances of death and no adverse effects on the liver and liver function (9,12). Longer toxicity analyses have not previously been performed, while only one long-term study of the glycemic effect of SC was identified in the literature (12). Thus, it is important to determine the glycemic effect of SC leaves over shorter periods (i.e., a maximum of 14 days).

The present study sought to determine the effect of SC leaf extract on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats after 14 days of treatment.

Materials and methods

Plant material and sample preparation

The SC leaves used in this study were obtained from Beringharjo Market, Yogyakarta, Indonesia. The leaf extract was made by macerating 3.2 g or 16.8 g of dried leaves in 100 mL of water for 24 hours. The liquid part was then separated from the residue by means of decantation.

Experimental animals

The protocol for this study was approved by the Health Research Ethics Committee of Universitas Alma Ata (Ref No: KE/AA/V/600/EC/2018). Male *Rattus norvegicus* rats weighing 185–224 g were used in this study. The rats were housed individually in an ambient condition with a 12 h light–dark cycle. The rats were fed AIN-93 diets (13) and given free access to drinking water. The rats' food intake and BW were recorded during the study.

Induction of diabetes and treatment

Thirty rats were divided into five groups, namely the normal/non-diabetic control group (negative control [NC] group), STZ-induced diabetic control group (positive control [PC] group), STZ-induced diabetic treated with 5 mg/kg BW glibenclamide group (GB group), STZ-induced diabetic treated with 3.2% SC leaf extract group (SC1 group), and STZ-induced diabetic treated with 16.8% SC leaf extract group (SC2 group). The positive control and diabetic treated groups were intraperitoneally injected with 110 mg/kg BW nicotinamide dissolved in 0.9% NaCl buffer saline prior to the injection of a single dose of 45 mg/kg BW STZ dissolved in citrate buffer to induce diabetes (14). The rats' diabetes status was determined on the third day following the STZ injection. All samples were administered orally as a single dose each day throughout the 14-day treatment period.

Blood sampling and biochemical analysis

Blood samples were collected from the rats' retroorbital vein by means of the microcapillary technique and then centrifuged at 400 rpm for 15 minutes to obtain the plasma. The rats' blood glucose levels and lipid profiles were measured three times, namely before and three days after the induction of diabetes as well as at the end of the intervention period. Rats with blood glucose levels up to 200 mg/dL were included in the diabetic groups. The utilized enzymatic assessment methods were glucose oxidase–peroxidase aminoantipyrin (GOD-PAP) for the glucose levels, glycerol phosphate oxidase–phenyl aminophyrazolon (GPO-PAP) for the triglyceride levels, and cholesterol oxidase–peroxidase aminoantipyrin (CHOD-PAP) for the cholesterol levels (total cholesterol, high density lipoprotein [HDL], and low-density lipoprotein [LDL]).

Statistical analysis

Data were performed as mean±standard deviation (SD) and analyzed by one-way analysis of variance (ANOVA) continued with Duncan's Multiple Range Test (DMRT) at $p < 0.05$. The analysis used statistical package for the social sciences (SPSS) software (version 16.0 SPSS Inc., Chicago, USA).

Results

Feed intake and body weight of rat

Feed intake of rats during 14 days of treatment tended to be stable and began to increase at the end period of the study (Figure 1). All treatment groups showed an increase in body weight, except PC (Figure 2), This indicated that the treatment of GB, SC1, and SC2 were able to reduce the effect of weight loss experienced by diabetic rats (PC).




Figure 1: Feed intake of rats during 14 days of intervention. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* leaf extract/SC1,16.8% SC2)




Figure 2: Bodyweight of rats during 14 days of intervention. Control groups (non- diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* leaf extract/SC1,16.8% SC2)

Effect of SC on blood glucose levels

Table 1: Effect of various treatments on blood glucose levels of rats

Groups	Blood glucose levels (mg/dL)			ΔK
	Before STZ	After STZ	After intervention (14	
	induction	induction (0 d)	d)	
NC	68.5±2.9 ^a	69.8±2.3 ^a	71.3±2.1*	+1.50 ^b
PC	68.7±3.0 ^a	260.0±5.0 ^b	260.9±5.0*	+0.83 ^b
GB	69.9±2.6 ^a	254.7±5.7 ^b	97.0±1.3*	-157.66 ^a
SC1	68.4±2.1 ^a	256.5±5.9 ^b	133.6±3.5*	-122.93 ^a
SC2	69.8±2.6 ^a	257.6±3.6 ^b	109.0±1.4*	-148.6 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1, 16.8% SC2)

It can be seen from Table 1 that the initial blood glucose level of rats was normal, at 68.4-69.9 mg/dL ($p > 0.05$). Except for NC, almost all treatment groups were given STZ to significantly increase blood glucose levels ($p < 0.05$). After 14 d of treatment, the blood glucose levels of GB, SC1, and SC2 groups were significantly lower than that of day 0 ($p < 0.05$). At the same time, in the PC (positive control) group, blood glucose levels increased. It was proved that SC has the same effect as the commercially available drug group, and the effective lowering level of blood glucose levels was 3.2% for 14 d of treatment.

Effect of SC on lipid profile

Before STZ administration, the plasma triglyceride levels of rats in all treatment groups were normal, between 74.5-76.6 mg/dL. Administering STZ to the diabetic group (PC, GB, SC1, and SC2) had an impact in increasing triglyceride levels in the range of 126.6-129.4 mg/dL. The GB and SC2 interventions were able to significantly reduced blood triglyceride levels with a decrease of 29.23 and 20.45 mg/dL, respectively ($p < 0.05$) (Table 2).

Table 2: Effect of various treatments on triglyceride levels of rats

Groups	Trygliceride levels (mg/dL)			ΔK
	Before STZ	After STZ	Af t er i n t e r v e n t i o n (1 4 d)	
	induct io n	induct io n (0 d)		
NC	76.6±2.5 ^a	78.2±2.0 ^a	81.9±1.9*	+3.74 ^d
PC	75.2±2.7 ^a	126.6±3.6 ^b	127.9±3.6*	+1.33 ^d
GB	74.5±2.0 ^a	127.0±3.6 ^b	97.9±2.5*	-29.23 ^a
SC1	74.6±2.5 ^a	129.4±1.5 ^b	127.6±2.2	-1.86 ^c
SC2	76.0±3.1 ^a	128.4±3.2 ^b	107.1±2.5*	-20.45 ^b

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

Table 3 shows that the total cholesterol levels in all treatment groups were the same, which was between 94-95.4 mg/dL. The presence of STZ induction in the diabetic rat groups significantly increased total cholesterol levels (p<0.05). SC was equivalent to GB in reducing the level of total cholesterol, the best concentration was 16.8% (SC2 group) (p<0.05).

Table 3: Effect of various treatments on total cholesterol levels of rats

Groups	Total cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction	After intervention	
		(0 d)	(14 d)	
NC	94.0±2.9 ^a	94.9±2.6 ^a	95.7±2.7	+0.75 ^c
PC	94.4±1.4 ^a	170.5±3.5 ^b	174.4±4.2	+3.86 ^c
GB	95.4±2.2 ^a	177.5±5.1 ^c	128.5±2.1*	-49.07 ^a
SC1	95.2±2.7 ^a	181.0±5.2 ^c	156.8±1.87*	-24.20 ^b
SC2	94.4±3.6 ^a	180.2±7.7 ^c	138.4±2.6*	-41.72 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

LDL cholesterol levels were in the range of 22.4-26.3 mg/dL which belonged to normal category. After being induced, LDL levels in diabetic rats increased and were significantly different compared to the normal rats' group (NC) (p<0.05). The 14-day intervention in the treatment group significantly reduced LDL cholesterol levels (p<0.05), while the untreated diabetes group (PC) did not. SC2 group had the highest reduction of cholesterol level with the same result with GB in the end of study (Table 4).

Table 4: Effect of various treatments on LDL cholesterol levels of rats

Groups	LDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 d)	After intervention (14 d)	
NC	25.7±1.13 ^b	25.2±1.12 ^a	27.5±1.28*	+6.9 ^c
PC	26.3 ±1.09 ^b	73.4±2.56 ^c	74.5±1.71	+1.1 ^c
GB	25.2 ±1.43 ^b	70.2±2.09 ^b	33.4±1.55*	-36.8 ^a
SC1	24.4 ±1.02 ^a	75.7±1.75 ^c	51.9±1.55*	-23.7 ^b
SC2	22.4±2.47 ^a	75.3±1.46 ^c	38.4±3.11*	-36.9 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% *S.crispus* extract/SC2)

The HDL cholesterol levels of all groups were classified as normal rats at the beginning of the study and ranged from 80.8-82 mg/dL. STZ induced a reduction in HDL levels. Treatments for 14 d increased HDL levels significantly (p<0.05) with the same increasing effect in all groups of rats (p>0.05). Administering SC at a minimum level (3.2%) was able to give the effect of increasing HDL levels which was equivalent to the commercial drug group (GB) (Table 5).

Table 5: Effect of various treatments on HDL cholesterol levels of rats

Groups	HDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 d)	After intervention (14 d)	
NC	81.9±1.41 ^a	80.1±1.33 ^b	79.5±0.99	+1.10 ^a
PC	81.6±1.07 ^a	25.3±1.43 ^a	25.1±1.20	-0.14 ^a
GB	82.0±1.41 ^a	25.1±0.38 ^a	68.5±1.46*	36.01 ^b
SC1	80.8±2.17 ^a	25.3±0.79 ^a	49.4±1.82*	24.14 ^b
SC2	81.6±1.55 ^a	24.1±1.62 ^a	62.5±2.08*	24.90 ^b

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

Discussion

Feed intake and bodyweight of rat

In this study, SC treatment did not affect the rats' feed intake, which proved that there was no effect of SC administration on the rats' appetite. During the 14 days of treatment, the weight of rats that received SC intervention also increased a lot, while the weight of untreated diabetic rats (PC) continued to drop. Weight loss in diabetic rats was caused by

continual fat catabolism to meet energy needs in their body as a result of the inability of cells to metabolize glucose in the blood. This inability of glucose metabolism could be caused by a disruption of pancreatic β -cells in producing insulin (15).

Effect of SC on blood glucose levels

The blood glucose levels of rats at the beginning of the study were categorized as normal, then significantly increased to >126 mg/dL or classified as high (16) after STZ induction. STZ induced DM by donating NO (nitric oxide) which contributed to pancreatic cell damage by increasing the activity of guanyl cyclase and formation of cGMP. NO was produced when STZ led to metabolism in cells. In addition, STZ was also able to generate reactive oxygen, which had a high role in pancreatic cell damage (17). In a previous study, the inducement of diabetic by STZ influenced disorientation of liver cellular structures, including glycogen deposition, nuclear location on the periphery of cell membranes, and acidophilic cytoplasm. Liver cell also experienced to necroticization, microvascular vacuolization, and hydropic inflammation (18).

In this study, indications of DM in rats could also be observed from the higher urinary frequency (polyuria). In general, people with diabetes have symptoms of polyuria (a lot of urination), polydipsia (a lot of drinking) and polyphagia (a lot of eating) with weight loss (19). SC treatment was able to significantly reduced blood glucose levels in rats and was comparable to the drug (GB group) ($p < 0.05$). These results were in line with previous study which showed that SC was effective in reducing blood glucose levels by giving the treatment at the dose of 14.7 mg/30g BW (20). In another study, processing SC into fermented tea also showed a decrease in blood glucose levels (10). This could be affected by the flavonoid content in SC which was able to act as an antioxidant by protecting pancreatic cell damage from the effects of free radicals, resulting in higher insulin sensitivity (21)(22). In addition, the antioxidant had the role in accelerating the release of glucose from the circulation which can reduce blood glucose levels (23).

Effect of SC on lipid profile

The lipid profile of rat at the beginning study including triglyceride, HDL, and LDL levels was normal (24)(25)(26), except total cholesterol level which was classified as high because the level was above 10-54 mg/dL (27). The presence of STZ induction changed the lipid profile to abnormal. As previously discussed, STZ was able to induce an increase in blood glucose levels (Table 1). It could be caused by insulin resistance affecting its fat metabolism (28). As a result of insulin resistance, sensitive hormone in adipose tissue increased, generated higher lipolysis of triglycerides in adipose tissue, leading to enhancement of free fatty acid levels in the blood. The excess of free fatty acids were converted into phospholipids, cholesterol, and some were used as raw materials for triglycerides generation in the liver which can then be transported into the blood in the form of lipoproteins (29). In this condition, abnormalities of fat metabolism occurred in the form of increased triglyceride levels, decreased HDL, increased small dense LDL subtraction, known as the atherogenic lipoprotein phenotype or lipid triad (30). Therefore,

the increase of blood glucose levels in DM patients had a significant correlation with the increase in triglyceride and cholesterol levels.

Administering SC for 14 days had a significant effect in improving lipid profiles. This can be seen from the SC's ability to significantly reduce triglyceride levels at a dose of 16.8% (SC2 group) and total cholesterol or LDL levels at a dose of 3.2% (SC1 group) ($p < 0.05$). However, by administering SC at a dose of 16.8%, the best reduction in total cholesterol and LDL cholesterol levels was achieved or equivalent to commercial drugs. HDL cholesterol levels can be increased to normal and equivalent to commercial drugs starting from the SC dose of 3.2% ($p < 0.05$).

This study was supported by previous experiment that using SC in the form of fermented tea which also showed improvements in lipid profiles such as lowering cholesterol, triglyceride, and LDL levels, but increasing HDL levels. In the form of SC ethanol extract at a dose of 400 mg/kg BW for 21 days of treatment, there was also a decrease in cholesterol and LDL levels (11). SC contains bioactive compounds such as alkaloids, saponins, β -sitosterol, flavonoids, tannins, potassium, and polyphenols. β -sitosterol was a hypocholesterolemic substance that might reduce cholesterol levels in the blood (10). The presence of β -sitosterol and flavonoids controlled cholesterol synthesis by inhibiting the activity of the enzyme acyl-CoA cholesterol acyl transferase (ACAT) in HepG2 cells which play a role in reducing cholesterol esterification in the intestine and liver. It also inhibited the activity of the HMG-CoA enzyme leading to inhibition of cholesterol synthesis (31). Flavonoids in SC reduced blood cholesterol levels by reducing or inhibiting the absorption of bile acids and cholesterol in the small intestine. It resulted in high excretion through feces leading to the formation of bile acids from cholesterol of liver cells. Increased excretion of fecal cholesterol also occurred because of the binding in the intestine between fiber with cholesterol and bile acids which will eventually be excreted or excreted through the feces. This reduced enterohepatic circulation of bile acids and increase the conversion of cholesterol into bile acids leading to the decrease of plasma cholesterol levels (32).

Conclusions

This study proved that SC extract provided good hypoglycemic effect starting at a concentration of 3.2%. However, administration of 16.8% SC was optimal not only in lowering blood glucose levels, but improving lipid profiles in the form of decreasing levels of triglycerides, total cholesterol, LDL cholesterol, and increasing HDL cholesterol.

Acknowledgments


The authors would like to thank Alma Ata University for the many facilities for this study.

References

1. Dewi PS, Anisa IN. Pengaruh Pemberian Ekstrak Etanol Sirih Merah Terhadap Kadar Glukosa Darah Pada Tikus Wistar Jantan Model Hiperkolesterolemia. *Kartika J Ilm Farm*. 2014;2(2).
2. Yahya J, Stephanie, Wijaya E. Pemanfaatan Teh Daun Keji Beling (*Strobilanthes Crispus* BL) sebagai Alternatif Pengobatan Penyakit Batu Ginjal dan Diabetes. Bogor; 2009.
3. Aprilia V, Kurniasari Y, Hadi H, Nurinda E, Ariftiyana S. Efek pemberian tepung porang (*Amorphophallus oncophyllus*) terhadap kadar glukosa darah, trigliserida, HDL, LDL, dan kolseterol darah tikus diabetes yang diinduksi streptozotosin. 2017.
4. Afifah E. Pemberian ekstrak air buah sawo (*Manilkara zapota* L.) menurunkan kadar glukosa darah tikus (*rattus norvegicus*) diabetes mellitus. *J Gizi dan Diet Indones (Indonesian J Nutr Diet*. 2016;3(3):180.
5. Francisco BM, Salvador M, Amparo N. Oxidative stress in myopia. *Oxid Med Cell Longev*. 2015;2015(3):225–36.
6. Purwaningtyastuti R, Nurwanti E, Huda N. Asupan vitamin C berhubungan dengan kadar glukosa darah pada pasien rawat jalan DM tipe 2. *J Gizi dan Diet Indones (Indonesian J Nutr Diet*. 2018;5(1):44.
7. Silalahi M. Pemanfaatan Kecibeling (*Strobilanthes crispa*) Sebagai Obat Tradisional dan Bioaktivitasnya. *J Edukasi Mat dan Sains*. 2020;IX:196–205.
8. Nurraihana H, Norfarizan-Hanoon NA. Phytochemistry, pharmacology and toxicology properties of *Strobilanthes crispus*. *Int Food Res J*. 2013;20(5):2045–56.
9. Norfarizan-Hanoon NA, Asmah R, Rokiah MY, Fauziah O, Faridah H. Absence of toxicity of *Strobilanthes crispa* juice in acute oral toxicity study in Sprague Dawley rats. *Sains Malaysiana*. 2012;41(4):403–9.
10. Fadzelly ABM, Asmah R, Fauziah O. Effects of *Strobilanthes crispus* tea aqueous extracts on glucose and lipid profile in normal and streptozotocin-induced hyperglycemic rats. *Plant Foods Hum Nutr*. 2006;61(1):7–12.
11. Oktavia S, Eriadi A, Valdis S. Pengaruh Pemberian Ekstrak Etanol Daun Kejibeling (*Strobilanthes Crispa* Blume) Terhadap Kadar Kolesterol Total Dan LDL Mencit Putih Kantan Hiperkolesterolemia. *J Farm Higea*. 2018;10(2):110–5.
12. Lim KT, Lim V, Chin JH. Subacute oral toxicity study of ethanolic leaves extracts of *Strobilanthes crispus* in rats. *Asian Pac J Trop Biomed*. 2012;2(12):948–52.
13. Reeves PG, Neilsen FH, Fahey, G. C J. AIN-93 purified diets for laboratory rodents: Final Report of the American Institute of Nutrition Ad Hoc Writing Committee on the Formulation of the AIN-76A Rodent Diet. *J Nutr*. 1993;123(11):1939–51.
14. Ghasemi A, Khalifi S, Jedi S. Streptozotocin-nicotinamide-induced rat model of type 2 diabetes (review). *Acta Physiol Hung*. 2014;101(4):408–20.
15. Rias YR, Sutikno E. Hubungan Antara Berat Badan Dengan Kadar Gula Darah

- Acak Pada Tikus Diabetes Mellitus. *J Wiyata*. 2017;4(1):72–7.
16. Association AD. Classification and Diagnosis of Diabetes Melitus. *Diabetes Care*. 2015;38.
 17. Nugroho AE. Animal Models Of Diabetes Mellitus : Pathology And Mechanism Of Some Diabetogenics. *Biodiversitas J Biol Divers*. 2006;7(4):378–82.
 18. Jaiswal YS, Tatke PA, Gabhe SY, Vaidya AB. Antidiabetic activity of extracts of *Anacardium occidentale* Linn. leaves on n-streptozotocin diabetic rats. *J Tradit Complement Med* [Internet]. 2016; Available from: <http://linkinghub.elsevier.com/retrieve/pii/S2225411016302899>
 19. Snehalata C, Ramachandran A. *Diabetes Melitus Dalam Gizi Kesehatan Masyarakat*. Jakarta, Indonesia: Penerbit Buku Kedokteran EGC; 2009.
 20. Nonci FY, Leboe DW, Armalia. Uji aktivitas ekstrak etanl daun kejjibeling (*Strobilanthes crispus* Linn) terhadap penurunan kadar glukosa darah pada mencit jantan (*Mus musculus*). *JF FIK UINAM*. 2016;4(1):16–20.
 21. Ismawati I, Sembirin LP. The Effect of Alpha Lipoic Acid in Preventing Atherosclerosis Lesion on the Abdominal Aorta of Diabetes Mellitus Type 2 Rats. *J Kedokt dan Kesehat*. 2020;16(1).
 22. Ashari A, Nurinda E, Fatmawati A. Peningkatan kadar insulin tikus wistar (*Rattus norvegicus*) jantan yang diinduksi streptozotosin akibat pemberian ekstrak etanol batang brotowali (*Tinospora crispa* L.). *INPHARNMED J (Indonesian Pharm Nat Med Journal)*. 2021;5(1):1.
 23. Widowati W. Potensi antioksidan sebagai antidiabetes. *J Kesehat Masy*. 2008;7(2):193–202.
 24. Suryaatmadja M. Pemeriksaan Trigliserida Tanpa Puasa. *Summit Lipid Update*. 2012;7:1–2.
 25. Nugraheni M, Lastariwati B, Purwanti S. Proximate and chemical analysis of gluten-free enriched, resistant starch type 3 from *Maranta arundinacea* flour and its potential as a functional food. *Pakistan J Nutr*. 2017;16(5):322–30.
 26. Adhi Pradana D, Dwiratna DW, Widyarini S. Aktivitas Ekstrak Etanolik Bayam Merah (*Amaranthus tricolor* L.) Terstandar sebagai Upaya Preventif Steatosis: Studi in Vivo. *J Sains Farm Klin*. 2017;3(2):120.
 27. Nurhidajah, Astuti R, Nurrahman. Black rice potential in HDL and LDL profile in sprague dawley rat with high cholesterol diet. *IOP Conf Ser Earth Environ Sci*. 2019;292(1).
 28. Trivedi N a, Mazumdar B, Bhatt JD, Hemavathi KG. Effect of shilajit on blood glucose and lipid profile in alloxan- induced diabetic rats. *Indian J Pharmacol*. 2004;36(6):373–6.
 29. Yassin MM, Mwafy SN. Protective potential of glimepiride and Nerium oleander extract on lipid profile, body growth rate, and renal function in streptozotocin-induced diabetic rats. *Turkish J Biol*. 2007;31(2):95–102.
 30. Adam J. *Dislipidemia: Ilmu Penyakit Dalam*. 4th ed. Jakarta, Indonesia: Pusat Penerbitan Departemen Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Indonesia; 2006.

31. Riyanto S, Muwarni R H. Yoghurt kedelai hitam (black soyghurt) dapat menurunkan kadar LDL tikus hiperkolesterolemia. *J Gizi dan Diet Indones* (Indonesian J Nutr Diet. 2016;3(1):1.
32. Potter SM. Overview of Proposed Mechanisms for the Hypocholesterolemic effects os Soy. *J Nutr*. 1995;125:606.

UNIVERSITAS
Alma Ata
Al-Furqan

Pemberitahuan Hasil Review: 21 Maret 2022

verianiaprilia verianiaprilia <verianiaprilia@almaata.ac.id>

[OAMJMS] Editor Decision

Ana Vucurevic via SFS - Journals (Scientific Foundation SPIROSKI - Journals), Skopje, Republic of Macedonia <noreply@publicknowledgeproject.org> 21 Maret 2022
pukul 20.08

Balas Ke: Ana Vucurevic <anavucurevic.nfs@gmail.com>

Kepada: Laras Ayu Fauziah Nur Fitri <larasayufauziah@gmail.com>, Tri Hernawati <Trihernawati87@gmail.com>, Desta Sari D <destasari30@gmail.com>, Indra Kusuma W <26@gmail.com>, Kunti Septiani <kuntiseptiani14@gmail.com>, Hamam Hadi <hhadi@almaata.ac.id>, Yulinda Kurniasari <kurniasari.yulinda@almaata.ac.id>, Effatul Afifah <effatulafifah@almaata.ac.id>, Veriani Aprilia <verianiaprilia@almaata.ac.id>

Laras Ayu Fauziah Nur Fitri, Tri Hernawati, Desta Sari D, Indra Kusuma W, Kunti Septiani, Hamam Hadi, Yulinda Kurniasari, Effatul Afifah, Veriani Aprilia (Author):

We have reached a decision regarding your submission to Open Access Macedonian Journal of Medical Sciences, "The effect of Strobilanthes crispus on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats ", Manuscript ID = OJS9468.

Our decision is to:

The effect of *Strobilanthes crispus* on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats

Laras Ayu Fauziah Nur Fitri¹, Tri Hernawati¹, Desta Sari D¹, Indra Kusuma W¹, Kunti S¹, Hamam Hadi², Yulinda Kurniasari¹, Effatul Afifah¹, Veriani Aprilia¹

¹*Department of Nutrition, Faculty of Health Sciences, Alma Ata University, Jl. Brawijaya 99 Tamantirto, Kasihan, Bantul, Yogyakarta 55183*

²*Graduate School of Public Health, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta 55183, Indonesia*

Corresponding author: Veriani Aprilia, *Department of Nutrition, Faculty of Health Sciences, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta 55183, Indonesia.*

E-mail: verianiaprilia@almaata.ac.id. ORCID iD: 0000-0002-6231-6487

Short title: *Strobilanthes crispus* effect on blood glucose levels and lipid profile

Funding: This research was supported by Universitas Alma Ata

The effect of *Strobilanthes crispus* on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats

Abstract

BACKGROUND: Diabetes mellitus is a metabolic disease associated with unhealthy eating patterns, a lack of physical activity, obesity, smoking habits, and family history. *Strobilanthes crispus* (SC) contains antioxidant compounds known to exert a hypoglycemic effect during long-term interventions in non-diabetic samples.

AIMS: This study sought to determine the effect of administering the extract of SC leaves on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats.

METHODS: A total of 30 male *Rattus norvegicus* rats were divided into control and treatment groups. The control groups consisted of negative (normal rats) and positive controls (diabetic rats), whereas treatment groups consisted of diabetic rats administered with 3.2% and 16.8% SC leaf extract, and glibenclamide as the drug comparison groups. Blood samples were collected from the rats' retroorbital vein for the purposes of glucose level and lipid profile measurement before and after the induction of diabetes with STZ as well as 14 days after the intervention. The collected data were analyzed statistically by means of a one-way analysis of variance (ANOVA) continued with Duncan's multiple range test.

RESULTS: The results showed the rats to be hyperglycemic experienced changes in their lipid profiles following the induction of diabetes. The administration of 3.2% SC leaf extract for 14 days reduced the rats' blood glucose levels, while the effect of the 16.8% SC leaf extract was more pronounced in terms of reducing the rats' blood glucose levels and improving their lipid profiles (reducing the triglyceride, total cholesterol, and low-density lipoprotein cholesterol levels, while increasing the high-density lipoprotein cholesterol levels).

CONCLUSION: This study found 16.8% SC leaf extract to exert a good hypoglycemic effect and lead to an improvement in the lipid profiles in diabetic rats.

Keywords: *Diabetes mellitus, Strobilanthes crispus, blood sugar, lipid profile, rat*

Introduction

Diabetes mellitus (DM) comprises a group of metabolic diseases associated with defects in insulin secretion, insulin action, or a combination of the two [1]. DM is currently the fourth leading cause of death worldwide [2]. Among the different types of DM, type 2 DM is known to result from unhealthy eating habits, a lack of physical activity, obesity, smoking habits, and/or family history [3].

Hyperglycemia in patients with type 2 DM has been found to increase the formation of free radicals such as the superoxide, hydrogen peroxide, nitric oxide, and hydroxyl radicals [4]. Oxidative stress can result if the levels of free radicals exceed the

antioxidants levels in the body, resulting in damage to the cell organelles and enzymes as well as increased lipid peroxidation. Increased levels of fat in the blood (dyslipidemia) serve as an indicator of the occurrence of lipid peroxidation [5]. In addition, oxidative stress can also cause complications such as eye disorders, kidney damage, and nerve damage [6].

The consumption of antioxidants could inhibit the activity of free radicals. Keji beling (*Strobilanthes crispus* [SC]) is a local Indonesian plant species known for its bioactive properties, especially when it comes to treating diabetes and improving lipid profiles (i.e., polyphenols, alkaloids, coumarins, flavonoids, iridoids, triterpenes, and sterols) [7][8]. Several in vivo studies involving various doses and formulas of SC have demonstrated these beneficial effects. For example, a prior study found the ethanol extract of SC (a dose of 14.7 mg/g body weight [BW]) to be effective in terms of reducing the blood glucose levels in rats, although the results did not reveal a curative effect in DM subjects [9]. Other studies have examined the effect SC leaf juice administered at doses of 1 mL/kg BW, 1.5 mL/kg BW, and 2 mL/kg BW over a period of 30 days [9] [10] and in the form of ethanol extract administered at doses of 100 mg/kg BW, 200 mg/kg BW, and 400 mg/kg BW over a period of 21 days [11]. With regard to the safety of SC leaves, a toxicity analysis was only performed for 14 days of administration, although it revealed no instances of death and no adverse effects on the liver and liver function [9,12]. Longer toxicity analyses have not previously been performed, while only one long-term study of the glycemic effect of SC was identified in the literature [12]. Thus, it is important to determine the glycemic effect of SC leaves over shorter periods (i.e., a maximum of 14 days).

The present study sought to determine the effect of SC leaf extract on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats after 14 days of treatment.

Materials and methods

Plant material and sample preparation

The SC leaves used in this study were obtained from Beringharjo Market, Yogyakarta, Indonesia. The leaf extract was made by macerating 3.2 g or 16.8 g of dried leaves in 100 mL of water for 24 hours. The liquid part was then separated from the residue by means of decantation.

Experimental animals

The protocol for this study was approved by the Health Research Ethics Committee of Universitas Alma Ata (Ref No: KE/AA/V/600/EC/2018). Male *Rattus norvegicus* rats weighing 185–224 g were used in this study. The rats were housed individually in an ambient condition with a 12 h light–dark cycle. The rats were fed AIN-93 diets [13] and given free access to drinking water. The rats' food intake and BW were recorded during the study.

Induction of diabetes and treatment

Thirty rats were divided into five groups, namely the normal/non-diabetic control group (negative control [NC] group), STZ-induced diabetic control group (positive control [PC] group), STZ-induced diabetic treated with 5 mg/kg BW glibenclamide group (GB group), STZ-induced diabetic treated with 3.2% SC leaf extract group (SC1 group), and STZ-induced diabetic treated with 16.8% SC leaf extract group (SC2 group). The positive control and diabetic treated groups were intraperitoneally injected with 110 mg/kg BW nicotinamide dissolved in 0.9% NaCl buffer saline prior to the injection of a single dose of 45 mg/kg BW STZ dissolved in citrate buffer to induce diabetes [14]. The rats' diabetes status was determined on the third day following the STZ injection. All samples were administered orally as a single dose each day throughout the 14-day treatment period.

Blood sampling and biochemical analysis

Blood samples were collected from the rats' retroorbital vein by means of the microcapillary technique and then centrifuged at 400 rpm for 15 minutes to obtain the plasma. The rats' blood glucose levels and lipid profiles were measured three times, namely before and three days after the induction of diabetes as well as at the end of the intervention period. Rats with blood glucose levels up to 200 mg/dL were included in the diabetic groups. The utilized enzymatic assessment methods were glucose oxidase–peroxidase aminoantipyrin (GOD-PAP) for the glucose levels, glycerol phosphate oxidase–phenyl aminophyrazolon (GPO-PAP) for the triglyceride levels, and cholesterol oxidase–peroxidase aminoantipyrin (CHOD-PAP) for the cholesterol levels (total cholesterol, high density lipoprotein [HDL], and low-density lipoprotein [LDL]).

Statistical analysis

Data were performed as mean±standard deviation (SD) and analyzed by one-way analysis of variance (ANOVA) continued with Duncan's Multiple Range Test (DMRT) at $p < 0.05$. The analysis used statistical package for the social sciences (SPSS) software (version 16.0 SPSS Inc., Chicago, USA).

Results

Feed intake and body weight of rat

Feed intake of rats during 14 days of treatment tended to be stable and began to increase at the end period of the study (Figure 1). All treatment groups showed an increase in body weight, except PC (Figure 2), This indicated that the treatment of GB, SC1, and SC2 were able to reduce the effect of weight loss experienced by diabetic rats (PC).




Figure 1: Feed intake of rats during 14 days of intervention. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* leaf extract/SC1,16.8% SC2)




Figure 2: Bodyweight of rats during 14 days of intervention. Control groups (non- diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* leaf extract/SC1,16.8% SC2)

Effect of SC on blood glucose levels

Table 1: Effect of various treatments on blood glucose levels of rats

Groups	Blood glucose levels (mg/dL)			ΔK
	Before STZ	After STZ	After intervention (14	
	induction	induction (0 d)	d)	
NC	68.5±2.9 ^a	69.8±2.3 ^a	71.3±2.1*	+1.50 ^b
PC	68.7±3.0 ^a	260.0±5.0 ^b	260.9±5.0*	+0.83 ^b
GB	69.9±2.6 ^a	254.7±5.7 ^b	97.0±1.3*	-157.66 ^a
SC1	68.4±2.1 ^a	256.5±5.9 ^b	133.6±3.5*	-122.93 ^a
SC2	69.8±2.6 ^a	257.6±3.6 ^b	109.0±1.4*	-148.6 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1, 16.8% SC2)

It can be seen from Table 1 that the initial blood glucose level of rats was normal, at 68.4-69.9 mg/dL ($p > 0.05$). Except for NC, almost all treatment groups were given STZ to significantly increase blood glucose levels ($p < 0.05$). After 14 d of treatment, the blood glucose levels of GB, SC1, and SC2 groups were significantly lower than that of day 0 ($p < 0.05$). At the same time, in the PC (positive control) group, blood glucose levels increased. It was proved that SC has the same effect as the commercially available drug group, and the effective lowering level of blood glucose levels was 3.2% for 14 d of treatment.

Effect of SC on lipid profile

Before STZ administration, the plasma triglyceride levels of rats in all treatment groups were normal, between 74.5-76.6 mg/dL. Administering STZ to the diabetic group (PC, GB, SC1, and SC2) had an impact in increasing triglyceride levels in the range of 126.6-129.4 mg/dL. The GB and SC2 interventions were able to significantly reduced blood triglyceride levels with a decrease of 29.23 and 20.45 mg/dL, respectively ($p < 0.05$) (Table 2).

Table 2: Effect of various treatments on triglyceride levels of rats

Groups	Triglyceride levels (mg/dL)			ΔK
	Before STZ	After STZ	After intervention (14 d)	
	induction	induction (0 d)		
NC	76.6±2.5 ^a	78.2±2.0 ^a	81.9±1.9*	+3.74 ^d
PC	75.2±2.7 ^a	126.6±3.6 ^b	127.9±3.6*	+1.33 ^d
GB	74.5±2.0 ^a	127.0±3.6 ^b	97.9±2.5*	-29.23 ^a
SC1	74.6±2.5 ^a	129.4±1.5 ^b	127.6±2.2	-1.86 ^c
SC2	76.0±3.1 ^a	128.4±3.2 ^b	107.1±2.5*	-20.45 ^b

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

Table 3 shows that the total cholesterol levels in all treatment groups were the same, which was between 94-95.4 mg/dL. The presence of STZ induction in the diabetic rat groups significantly increased total cholesterol levels (p<0.05). SC was equivalent to GB in reducing the level of total cholesterol, the best concentration was 16.8% (SC2 group) (p<0.05).

Table 3: Effect of various treatments on total cholesterol levels of rats

Groups	Total cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction	After intervention	
		(0 d)	(14 d)	
NC	94.0±2.9 ^a	94.9±2.6 ^a	95.7±2.7	+0.75 ^c
PC	94.4±1.4 ^a	170.5±3.5 ^b	174.4±4.2	+3.86 ^c
GB	95.4±2.2 ^a	177.5±5.1 ^c	128.5±2.1*	-49.07 ^a
SC1	95.2±2.7 ^a	181.0±5.2 ^c	156.8±1.87*	-24.20 ^b
SC2	94.4±3.6 ^a	180.2±7.7 ^c	138.4±2.6*	-41.72 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

LDL cholesterol levels were in the range of 22.4-26.3 mg/dL which belonged to normal category. After being induced, LDL levels in diabetic rats increased and were significantly different compared to the normal rats' group (NC) (p<0.05). The 14-day intervention in the treatment group significantly reduced LDL cholesterol levels (p<0.05), while the untreated diabetes group (PC) did not. SC2 group had the highest reduction of cholesterol level with the same result with GB in the end of study (Table 4).

Table 4: Effect of various treatments on LDL cholesterol levels of rats

Groups	LDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 d)	After intervention (14 d)	
NC	25.7±1.13 ^b	25.2±1.12 ^a	27.5±1.28*	+6.9 ^c
PC	26.3 ±1.09 ^b	73.4±2.56 ^c	74.5±1.71	+1.1 ^c
GB	25.2 ±1.43 ^b	70.2±2.09 ^b	33.4±1.55*	-36.8 ^a
SC1	24.4 ±1.02 ^a	75.7±1.75 ^c	51.9±1.55*	-23.7 ^b
SC2	22.4±2.47 ^a	75.3±1.46 ^c	38.4±3.11*	-36.9 ^a

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% *S.crispus* extract/SC2)

The HDL cholesterol levels of all groups were classified as normal rats at the beginning of the study and ranged from 80.8-82 mg/dL. STZ induced a reduction in HDL levels. Treatments for 14 d increased HDL levels significantly (p<0.05) with the same increasing effect in all groups of rats (p>0.05). Administering SC at a minimum level (3.2%) was able to give the effect of increasing HDL levels which was equivalent to the commercial drug group (GB) (Table 5).

Table 5: Effect of various treatments on HDL cholesterol levels of rats

Groups	HDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 d)	After intervention (14 d)	
NC	81.9±1.41 ^a	80.1±1.33 ^b	79.5±0.99	+1.10 ^a
PC	81.6±1.07 ^a	25.3±1.43 ^a	25.1±1.20	-0.14 ^a
GB	82.0±1.41 ^a	25.1±0.38 ^a	68.5±1.46*	36.01 ^b
SC1	80.8±2.17 ^a	25.3±0.79 ^a	49.4±1.82*	24.14 ^b
SC2	81.6±1.55 ^a	24.1±1.62 ^a	62.5±2.08*	24.90 ^b

*) Significantly different between 0 d and 14 d using T-test. Superscript values in the same column are significantly different (p< 0.05) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values show the decrease and increase of blood glucose levels between 0 d and 14 d of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *S.crispus* extract/SC1,16.8% SC2)

Discussion

Feed intake and bodyweight of rat

In this study, SC treatment did not affect the rats' feed intake, which proved that there was no effect of SC administration on the rats' appetite. During the 14 days of treatment, the weight of rats that received SC intervention also increased a lot, while the weight of untreated diabetic rats (PC) continued to drop. Weight loss in diabetic rats was caused by

continual fat catabolism to meet energy needs in their body as a result of the inability of cells to metabolize glucose in the blood. This inability of glucose metabolism could be caused by a disruption of pancreatic β -cells in producing insulin [15].

Effect of SC on blood glucose levels

The blood glucose levels of rats at the beginning of the study were categorized as normal, then significantly increased to >126 mg/dL or classified as high [16] after STZ induction. STZ induced DM by donating NO (nitric oxide) which contributed to pancreatic cell damage by increasing the activity of guanyl cyclase and formation of cGMP. NO was produced when STZ led to metabolism in cells. In addition, STZ was also able to generate reactive oxygen, which had a high role in pancreatic cell damage [17]. In a previous study, the inducement of diabetic by STZ influenced disorientation of liver cellular structures, including glycogen deposition, nuclear location on the periphery of cell membranes, and acidophilic cytoplasm. Liver cell also experienced to necroticization, microvascular vacuolization, and hydropic inflammation [18].

In this study, indications of DM in rats could also be observed from the higher urinary frequency (polyuria). In general, people with diabetes have symptoms of polyuria (a lot of urination), polydipsia (a lot of drinking) and polyphagia (a lot of eating) with weight loss [19]. SC treatment was able to significantly reduced blood glucose levels in rats and was comparable to the drug (GB group) ($p < 0.05$). These results were in line with previous study which showed that SC was effective in reducing blood glucose levels by giving the treatment at the dose of 14.7 mg/30g BW [20]. In another study, processing SC into fermented tea also showed a decrease in blood glucose levels [10]. This could be affected by the flavonoid content in SC which was able to act as an antioxidant by protecting pancreatic cell damage from the effects of free radicals, resulting in higher insulin sensitivity [21][22]. In addition, the antioxidant had the role in accelerating the release of glucose from the circulation which can reduce blood glucose levels [23].

Effect of SC on lipid profile

The lipid profile of rat at the beginning study including triglyceride, HDL, and LDL levels was normal [24][25][26], except total cholesterol level which was classified as high because the level was above 10-54 mg/dL [27]. The presence of STZ induction changed the lipid profile to abnormal. As previously discussed, STZ was able to induce an increase in blood glucose levels (Table 1). It could be caused by insulin resistance affecting its fat metabolism [28]. As a result of insulin resistance, sensitive hormone in adipose tissue increased, generated higher lipolysis of triglycerides in adipose tissue, leading to enhancement of free fatty acid levels in the blood. The excess of free fatty acids were converted into phospholipids, cholesterol, and some were used as raw materials for triglycerides generation in the liver which can then be transported into the blood in the form of lipoproteins [29]. In this condition, abnormalities of fat metabolism occurred in the form of increased triglyceride levels, decreased HDL, increased small dense LDL subtraction, known as the atherogenic lipoprotein phenotype or lipid triad [30]. Therefore,

the increase of blood glucose levels in DM patients had a significant correlation with the increase in triglyceride and cholesterol levels.

Administering SC for 14 days had a significant effect in improving lipid profiles. This can be seen from the SC's ability to significantly reduce triglyceride levels at a dose of 16.8% (SC2 group) and total cholesterol or LDL levels at a dose of 3.2% (SC1 group) ($p < 0.05$). However, by administering SC at a dose of 16.8%, the best reduction in total cholesterol and LDL cholesterol levels was achieved or equivalent to commercial drugs. HDL cholesterol levels can be increased to normal and equivalent to commercial drugs starting from the SC dose of 3.2% ($p < 0.05$).

This study was supported by previous experiment that using SC in the form of fermented tea which also showed improvements in lipid profiles such as lowering cholesterol, triglyceride, and LDL levels, but increasing HDL levels. In the form of SC ethanol extract at a dose of 400 mg/kg BW for 21 days of treatment, there was also a decrease in cholesterol and LDL levels [11]. SC contains bioactive compounds such as alkaloids, saponins, β -sitosterol, flavonoids, tannins, potassium, and polyphenols. β -sitosterol was a hypocholesterolemic substance that might reduce cholesterol levels in the blood [10]. The presence of β -sitosterol and flavonoids controlled cholesterol synthesis by inhibiting the activity of the enzyme acyl-CoA cholesterol acyl transferase (ACAT) in HepG2 cells which play a role in reducing cholesterol esterification in the intestine and liver. It also inhibited the activity of the HMG-CoA enzyme leading to inhibition of cholesterol synthesis [31]. Flavonoids in SC reduced blood cholesterol levels by reducing or inhibiting the absorption of bile acids and cholesterol in the small intestine. It resulted in high excretion through feces leading to the formation of bile acids from cholesterol of liver cells. Increased excretion of fecal cholesterol also occurred because of the binding in the intestine between fiber with cholesterol and bile acids which will eventually be excreted or excreted through the feces. This reduced enterohepatic circulation of bile acids and increase the conversion of cholesterol into bile acids leading to the decrease of plasma cholesterol levels [32].

Conclusions

This study proved that SC extract provided good hypoglycemic effect starting at a concentration of 3.2%. However, administration of 16.8% SC was optimal not only in lowering blood glucose levels, but improving lipid profiles in the form of decreasing levels of triglycerides, total cholesterol, LDL cholesterol, and increasing HDL cholesterol.

Acknowledgments

The authors would like to thank Alma Ata University for the many facilities for this study.

References

1. Dewi PS, Anisa IN. Effect of ethanolic extract of red betel on glucose level of hypercholesterolemic male Wistar rat. *Kartika J Ilm Farm*. 2014;2(2). (Indonesian).
2. Yahya J, Stephanie, Wijaya E. Utilization of keji beling (*Strobilanthes Crispus* BL) tea as alternative medicine of kidney stones and diabetes. Bogor; 2009. (Indonesian).
3. Aprilia V, Kurniasari Y, Hadi H, Nurinda E, Ariftiyana S. Effect of porang flour (*Amorphophallus oncophyllus*) on blood glucose, triglyceride, HDL, LDL, and cholesterol of Streptozotocin-induced diabetic. 2017. (Indonesian).
4. Afifah E. Addition of sapodilla fruit (*Manilkara zapota* L.) decreased the blood glucose levels of diabetic rat (*rattus norvegicus*). *J Gizi dan Diet Indones* (Indonesian *J Nutr Diet*. 2016;3(3):180. (Indonesian)
5. Francisco BM, Salvador M, Amparo N. Oxidative stress in myopia. *Oxid Med Cell Longev*. 2015;2015(3):225–36.
6. Purwaningtyastuti R, Nurwanti E, Huda N. Vitamin C consumption related with blood glucose levels of type2-diabetes outpatients. *J Gizi dan Diet Indones* (Indonesian *J Nutr Diet*. 2018;5(1):44. (Indonesian).
7. Silalahi M. utilization of keji beling (*Strobilanthes crispa*) as traditional medicine and its bioactivity. *J Edukasi Mat dan Sains*. 2020;IX:196–205. (Indonesian).
8. Nurraihana H, Norfarizan-Hanoon NA. Phytochemistry, pharmacology and toxicology properties of *Strobilanthes crispus*. *Int Food Res J*. 2013;20(5):2045–56.
9. Norfarizan-Hanoon NA, Asmah R, Rokiah MY, Fauziah O, Faridah H. Absence of toxicity of *Strobilanthes crispa* juice in acute oral toxicity study in Sprague Dawley rats. *Sains Malaysiana*. 2012;41(4):403–9.
10. Fadzelly ABM, Asmah R, Fauziah O. Effects of *Strobilanthes crispus* tea aqueous extracts on glucose and lipid profile in normal and streptozotocin-induced hyperglycemic rats. *Plant Foods Hum Nutr*. 2006;61(1):7–12.
11. Oktavia S, Eriadi A, Valdis S. Effect of ethanolic extract of keji beling (*Strobilanthes crispa* Blume) leave on cholesterol levels and LDL of hypocholesterolemic male white rats. *J Farm Higea*. 2018;10(2):110–5. (Indonesian).
12. Lim KT, Lim V, Chin JH. Subacute oral toxicity study of ethanolic leaves extracts of *Strobilanthes crispus* in rats. *Asian Pac J Trop Biomed*. 2012;2(12):948–52.
13. Reeves PG, Neilsen FH, Fahey, G. C J. AIN-93 purified diets for laboratory rodents: Final Report of the American Institute of Nutrition Ad Hoc Writing Committee on the Formulation of the AIN-76A Rodent Diet. *J Nutr*. 1993;123(11):1939–51.
14. Ghasemi A, Khalifi S, Jedi S. Streptozotocin-nicotinamide-induced rat model of type 2 diabetes (review). *Acta Physiol Hung*. 2014;101(4):408–20.
15. Rias YR, Sutikno E. Relationship between body weight and random blood glucose levels of diabetic rats. *J Wiyata*. 2017;4(1):72–7. (Indonesian).

16. Association AD. Classification and Diagnosis of Diabetes Melitus. *Diabetes Care*. 2015;38.
17. Nugroho AE. Animal Models Of Diabetes Mellitus : Pathology And Mechanism Of Some Diabetogenics. *Biodiversitas J Biol Divers*. 2006;7(4):378–82.
18. Jaiswal YS, Tatke PA, Gabhe SY, Vaidya AB. Antidiabetic activity of extracts of *Anacardium occidentale* Linn. leaves on n-streptozotocin diabetic rats. *J Tradit Complement Med* [Internet]. 2016; Available from: <http://linkinghub.elsevier.com/retrieve/pii/S2225411016302899>
19. Snehalata C, Ramachandran A. Diabetes mellitus in public nutrition and health. Jakarta, Indonesia: Penerbit Buku Kedokteran EGC; 2009. (Indonesian).
20. Nonci FY, Leboe DW, Armalia. Activity of ethanolic extract of keji beling (*Strobilanthes crispus* Linn) in decreasing of blood glucose levels of male rats (*Mus musculus*). *JF FIK UINAM*. 2016;4(1):16–20. (Indonesian).
21. Ismawati I, Sembirin LP. The Effect of Alpha Lipoic Acid in Preventing Atherosclerosis Lesion on the Abdominal Aorta of Diabetes Mellitus Type 2 Rats. *J Kedokt dan Kesehat*. 2020;16(1).
22. Ashari A, Nurinda E, Fatmawati A. Increase of insulin levels of Streptozotocin-induced male Wistar rats (*Rattus norvegicus*) after ethanolic extract treatment of brotowali stem (*Tinospora crispa* L.). *INPHARNMED J (Indonesian Pharm Nat Med Journal)*. 2021;5(1):1. (Indonesian).
23. Widowati W. Potency of antioxidant as antidiabetes. *J Kesehat Masy*. 2008;7(2):193–202. (Indonesian).
24. Suryaatmadja M. Triglyceride test without fasting. *Summit Lipid Update*. 2012;7:1–2. (Indonesian).
25. Nugraheni M, Lastariwati B, Purwanti S. Proximate and chemical analysis of gluten-free enriched, resistant starch type 3 from *Maranta arundinacea* flour and its potential as a functional food. *Pakistan J Nutr*. 2017;16(5):322–30.
26. Adhi Pradana D, Dwiratna DW, Widyarini S. Activity of standardized-ethanolic extract of red spinach (*Amaranthus tricolor* L.) as preventive efforts of steatosis: In vivo study. *J Sains Farm Klin*. 2017;3(2):120. (Indonesian).
27. Nurhidajah, Astuti R, Nurrahman. Black rice potential in HDL and LDL profile in sprague dawley rat with high cholesterol diet. *IOP Conf Ser Earth Environ Sci*. 2019;292(1).
28. Trivedi N a, Mazumdar B, Bhatt JD, Hemavathi KG. Effect of shilajit on blood glucose and lipid profile in alloxan- induced diabetic rats. *Indian J Pharmacol*. 2004;36(6):373–6.
29. Yassin MM, Mwafy SN. Protective potential of glimepiride and Nerium oleander extract on lipid profile, body growth rate, and renal function in streptozotocin-induced diabetic rats. *Turkish J Biol*. 2007;31(2):95–102.
30. Adam J. *Dislipidemia: Internal medicine science*. 4th ed. Jakarta, Indonesia: Pusat Department of Internal Medicine Science, Medicine Faculty. Universitas Indonesia; 2006. (Indonesian).
31. Riyanto S, Muwarni R H. Black soyghurt lowered LDL levels of

- hypercholesterolemic rats. *J Gizi dan Diet Indones (Indonesian J Nutr Diet.* 2016;3(1):1. (Indonesian).
32. Potter SM. Overview of Proposed Mechanisms for the Hypocholesterolemic effects of Soy. *J Nutr.* 1995;125:606.



Proses produksi: 30 Maret 2022

verianiaprilia verianiaprilia <verianiaprilia@almaata.ac.id>

[OAMJMS] Editor Decision

Teodora Fildishevskia via SFS - Journals (Scientific Foundation SPIROSKI - Journals), Skopje, Republic of Macedonia <noreply@publicknowledgeproject.org>

30 Maret 2022
pukul 16.24

Balas Ke: Teodora Fildishevskia <tfildishevskia@id-press.eu>

Kepada: Laras Ayu Fauziah Nur Fitri <larasayufauziah@gmail.com>, Tri Hernawati <Trihernawati87@gmail.com>, Desta Sari D <destasari30@gmail.com>, Indra Kusuma W <26@gmail.com>, Kunti Septiani <kuntiseptiani14@gmail.com>, Hamam Hadi <hhadi@almaata.ac.id>, Yulinda Kurniasari <kurniasari.yulinda@almaata.ac.id>, Effatul Afifah <effatulafifah@almaata.ac.id>, Veriani Aprilia <verianiaprilia@almaata.ac.id>

Laras Ayu Fauziah Nur Fitri, Tri Hernawati, Desta Sari D, Indra Kusuma W, Kunti Septiani, Hamam Hadi, Yulinda Kurniasari, Effatul Afifah, Veriani Aprilia (Author):

The editing of your submission, "The effect of Strobilanthes crispus on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats ," Manuscript ID = OJS9468 is complete. We are now sending it to production.

Submission URL: <https://oamjms.eu/index.php/mjms/authorDashboard/submission/9468>

Copyright Form

Manuscript Number: OJS9468

This Agreement is made with *Open Access Macedonian Journal of Medical Sciences (OAMJMS)*. The Agreement is for the Article submitted by you ("Author") for publication *Open Access Macedonian Journal of Medical Sciences (OAMJMS)*. To enable publishing the Article in *Open Access Macedonian Journal of Medical Sciences (OAMJMS)* ("the Journal"), the ownership of copyright must be established. Please read and complete the form below (Section A plus Section B) and return one copy to the address given below. There are explanatory notes to this Agreement, which also form part of the Agreement.

The Article cannot be published until this signed Agreement is received by OAMJMS.

SECTION A: Author's Warranty (Please print your details)

Name: Veriani Aprilia

Address: Department of Nutrition, Faculty of Health Sciences, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta 55183, Indonesia.

Article title: The effect of *Strobilanthes crispus* on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats.

Names of all authors in the order in which they appear in the Article:

Laras Ayu Fauziah Nur Fitri, Tri Hernawati, Desta Sari D, Indra Kusuma W, Kunti S, Hamam Hadi, Yulinda Kurniasari, Effatul Afifah, Veriani Aprilia

In consideration of the publication of the Article in the above Journal, I hereby warrant and undertake:

- that this Article is an original work, has not been published before and is not being considered for publication elsewhere in its final form either in printed or electronic form.
- that I have obtained permission from the copyright holder to reproduce in the Article (in all media including print and electronic form) material not owned by me, and that I have acknowledged the source;
- that this Article contains no violation of any existing copyright or other third party right or any material of an obscene, indecent or otherwise unlawful nature and that to the best of my knowledge this Article does not infringe the rights of others;
- that I will indemnify and keep indemnified the Editors, Open Access Macedonian Journal of Medical Sciences against all claims and expenses (including legal costs and expenses) arising from any breach of this warranty and the other warranties on my behalf in this Agreement;
- that in the case of a multi-authored Article I have obtained copyright assignment from all co-authors, in writing, and authorization to enter into this Agreement on their behalf and that all co-authors have read and agreed the above warranties;



Author: Signed: Veriani Aprilia Date: March 31, 2022

SECTION B: Copyright Assignment

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Notwithstanding the above, I retain all proprietary rights other than copyright, such as patent and trade mark rights and rights to any process or procedure described in the Article.



Author: Signed: Veriani Aprilia Date: March 31, 2022

In signing this form, the signee asserts that any authors not signing have authorised the signee to do so on their behalf, and that the manuscript submitted has been approved by these authors in the form in which it has been submitted, and that the warranties given above have been read and agreed by all authors.

PRODUCTION Proof read: 20 Juli 2023

7/20/23, 1:04 PM

Gmail - Fwd: [OAMJMS] Proofreading Request (Author)



veriani aprilia <aprilia.triwanto@gmail.com>

Fwd: [OAMJMS] Proofreading Request (Author)

1 message

Laras Ayu Fauziah <larasayufauziah@gmail.com>
To: veriani aprilia <aprilia.triwanto@gmail.com>

Thu, Jul 20, 2023 at 10:25 AM

----- Forwarded message -----

From: **Teodora Fildishevska via SFS - Journals (Scientific Foundation SPIROSKI - Journals), Skopje, Republic of Macedonia** <noreply@publicknowledgeproject.org>
Date: Wed, 30 Mar 2022 at 16:24
Subject: [OAMJMS] Proofreading Request (Author)
To: Laras Ayu Fauziah Nur Fitri <larasayufauziah@gmail.com>

Dear Laras Ayu Fauziah Nur Fitri:

Your submission "The effect of Strobilanthes crispus on blood glucose levels and lipid profile of streptozotocin-induced diabetic rats " to Open Access Macedonian Journal of Medical Sciences now needs to be proofread by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and view PROOFING INSTRUCTIONS
3. Click on VIEW PROOF in Layout and proof the galley in the one or more formats used.
4. Enter corrections (typographical and format) in Proofreading Corrections using "Mark up text with edits" or "Use annotation and drawing markup tools to add comments in PDFs".
5. Save corrected Galley Proof with the addition "Corrected" to the file name (i.e., OAMJMS-9A-4528.pdf should be saved as OAMJMS-9A-4528-Corrected.pdf).
6. Click on the link "Add Discussion", define Subject "Corrected Galley Proof", write text in the field Message, Upload File in the field Attached Files.
7. Complete and sign **Copyright Form**, upload in the field Attached Files, and complete the message with clicking "OK".

Submission URL: <https://oamjms.eu/index.php/mjms/authorDashboard/submission/9468>
Username: Laras Ayu Fauziah Nur Fitri

Teodora Fildishevska
ID Design 2012/DOOEL Skopje, Skopje, Republic of Macedonia
tfildishevska@jd-press.eu

Workflow Publication

Submission Review Copyediting Production

Production Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
[OAMJMS] Proofreading Request (Author)	tfildishevska	-	0	<input type="checkbox"/>
▶ Corrected Galley Proof	lfitri	-	0	<input type="checkbox"/>
[OAMJMS] Your Article was Published	ivos	-	0	<input type="checkbox"/>

https://oamjms.eu/index.php/mjms/.../tab/author-dashboard/author-dashboard-tab/fetch-tab?submissionId=9468&stagedId=5

WhatsApp Image...jpeg

Show all X



The Effect of *Strobilanthes crispus* on Blood Glucose Levels and Lipid Profile of Streptozotocin-induced Diabetic Rats

Laras Ayu Fauziah Nur Fitri¹, Tri Hernawati¹, D. Desta Sari¹, W. Indra Kusuma¹, S. Kunti¹, Hamam Hadi², Yulinda Kurniasari¹, Effatul Afifah¹, Veriani Aprilia^{1*}

¹Department of Nutrition, Faculty of Health Sciences, Alma Ata University, Bantul, Yogyakarta, Indonesia; ²Graduate School of Public Health, Universitas Alma Ata, Bantul, Yogyakarta, Indonesia

Abstract

Edited by: <https://publons.com/researcher/3919877/mirko-spiroski>

Citation: Fitri LAFN, Hernawati T, Sari DD, Kusuma WI, Kunti S, Hadi H, Kurniasari Y, Afifah E, Aprilia V. The Effect of *Strobilanthes crispus* on Blood Glucose Levels and Lipid Profile of Streptozotocin-induced Diabetic Rats. Open-Access Maced J Med Sci. 2022 Feb 05; 10(T8):1-6. <https://doi.org/10.3889/oamjms.2022.9468>

Keywords: Diabetes mellitus; *Strobilanthes crispus*; Blood sugar; Lipid profile; Rat

*Correspondence: Veriani Aprilia, Department of Nutrition, Faculty of Health Sciences, Universitas Alma Ata, Jl. Brawijaya No. 99, Tamantirto, Kasihan, Bantul, Yogyakarta - 55183, Indonesia. E-mail: verianiaprilia@almaata.ac.id

Received: ???

Revised: ???

Accepted: ???

Copyright: © 2022 Laras Ayu Fauziah Nur Fitri, Tri Hernawati, D. Desta Sari, W. Indra Kusuma, S. Kunti, Hamam Hadi, Yulinda Kurniasari, Effatul Afifah, Veriani Aprilia

Funding: This research was supported by Universitas Alma Ata

Competing Interest: The authors have declared that no competing interest exists

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Diabetes mellitus is a metabolic disease associated with unhealthy eating patterns, a lack of physical activity, obesity, smoking habits, and family history. *Strobilanthes crispus* (SC) contains antioxidant compounds known to exert a hypoglycemic effect during long-term interventions in non-diabetic samples.

AIMS: This study sought to determine the effect of administering the extract of SC leaves on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats.

METHODS: A total of 30 male *Rattus norvegicus* rats were divided into control and treatment groups. The control groups consisted of negative (normal rats) and positive controls (diabetic rats), whereas the treatment groups consisted of diabetic rats administered with 3.2% and 16.8% SC leaf extract, and glibenclamide as the drug comparison groups. Blood samples were collected from the rats' retro-orbital vein for the purposes of glucose level and lipid profile measurement before and after the induction of diabetes with STZ as well as 14 days after the intervention. The collected data were analyzed statistically by means of a one-way analysis of variance continued with Duncan's multiple range test.

RESULTS: The results showed the rats to be hyperglycemic experienced changes in their lipid profiles following the induction of diabetes. The administration of 3.2% SC leaf extract for 14 days reduced the rats' blood glucose levels, while the effect of the 16.8% SC leaf extract was more pronounced in terms of reducing the rats' blood glucose levels and improving their lipid profiles (reducing the triglyceride, total cholesterol, and low-density lipoprotein cholesterol levels, while increasing the high-density lipoprotein cholesterol levels).

CONCLUSIONS: This study found 16.8% SC leaf extract to exert a good hypoglycemic effect and leads to an improvement in the lipid profiles in diabetic rats.

AQ1

Introduction

Diabetes mellitus (DM) comprises a group of metabolic diseases associated with defects in insulin secretion, insulin action, or a combination of the two [1]. DM is currently the fourth leading cause of death worldwide [2]. Among the different types of DM, type 2 DM is known to result from unhealthy eating habits, a lack of physical activity, obesity, smoking habits, and/or family history [3].

Hyperglycemia in patients with type 2 DM has been found to increase the formation of free radicals such as the superoxide, hydrogen peroxide, nitric oxide (NO), and hydroxyl radicals [4]. Oxidative stress can result if the levels of free radicals exceed the antioxidants levels in the body, resulting in damage to the cell organelles and enzymes as well as increased lipid peroxidation. Increased levels of fat in the blood (dyslipidemia) serve as an indicator of the occurrence of lipid peroxidation [5]. In addition, oxidative stress can

also cause complications such as eye disorders, kidney damage, and nerve damage [6].

The consumption of antioxidants could inhibit the activity of free radicals. Keji beling (*Strobilanthes crispus* [SC]) is a local Indonesian plant species known for its bioactive properties, especially when it comes to treating diabetes and improving lipid profiles (i.e., polyphenols, alkaloids, coumarins, flavonoids, iridoids, triterpenes, and sterols) [7], [8]. Several *in vivo* studies involving various doses and formulas of SC have demonstrated these beneficial effects. For example, a prior study found the ethanol extract of SC (a dose of 14.7 mg/g body weight [BW]) to be effective in terms of reducing the blood glucose levels in rats, although the results did not reveal a curative effect in DM subjects [9]. Other studies have examined the effect SC leaf juice administered at doses of 1 mL/kg BW, 1.5 mL/kg BW, and 2 mL/kg BW over a period of 30 days [9], [10] and in the form of ethanol extract administered at doses of 100 mg/kg BW, 200 mg/kg BW, and 400 mg/kg BW over a period of 21 days [11]. With regard to the safety of SC leaves, a toxicity analysis was

only performed for 14 days of administration, although it revealed no instances of death and no adverse effects on the liver and liver function [9], [12]. Longer toxicity analyses have not previously been performed, while only one long-term study of the glycemic effect of SC was identified in the literature [12]. Thus, it is important to determine the glycemic effect of SC leaves over shorter periods (i.e., a maximum of 14 days).

The present study sought to determine the effect of SC leaf extract on the blood glucose levels and blood serum lipid profiles of streptozotocin (STZ)-induced diabetic rats after 14 days of treatment.

as a single dose each day throughout the 14-day treatment period.

Methods

Plant material and sample preparation

The SC leaves used in this study were obtained from Beringharjo Market, Yogyakarta, Indonesia. The leaf extract was made by macerating 3.2 g or 16.8 g of dried leaves in 100 mL of water for 24 h. The liquid part was then separated from the residue by means of decantation.

Experimental animals

The protocol for this study was approved by the Health Research Ethics Committee of Universitas Alma Ata (Ref No: KE/AA/V/600/EC/2018). Male *Rattus norvegicus* rats weighing 185–224 g were used in this study. The rats were housed individually in an ambient condition with a 12 h light-dark cycle. The rats were fed AIN-93 diets [13] and given free access to drinking water. The rats' food intake and BW were recorded during the study.

Induction of diabetes and treatment

Thirty rats were divided into five groups, namely, the normal/non-diabetic control group (negative control [NC] group), STZ-induced diabetic control group (positive control [PC] group), STZ-induced diabetic treated with 5 mg/kg BW glibenclamide group (GB group), STZ-induced diabetic treated with 3.2% SC leaf extract group (SC1 group), and STZ-induced diabetic treated with 16.8% SC leaf extract group (SC2 group). The PC and diabetic-treated groups were intraperitoneally injected with 110 mg/kg BW nicotinamide dissolved in 0.9% NaCl buffer saline before the injection of a single dose of 45 mg/kg BW STZ dissolved in citrate buffer to induce diabetes [14]. The rats' diabetes status was determined on the 3rd day following the STZ injection. All samples were administered orally

Blood sampling and biochemical analysis

Blood samples were collected from the rats' retro-orbital vein by means of the microcapillary technique and then centrifuged at 400 rpm for 15 min to obtain the plasma. The rats' blood glucose levels and lipid profiles were measured three times, namely, before and 3 days after the induction of diabetes as well as at the end of the intervention period. Rats with blood glucose levels up to 200 mg/dL were included in the diabetic groups. The utilized enzymatic assessment methods were glucose oxidase- peroxidase aminoantipyrine for the glucose levels, glycerol phosphate oxidase-phenyl aminopyrazolone for the triglyceride levels, and cholesterol oxidase-peroxidase aminoantipyrine for the cholesterol levels (total cholesterol, high-density lipoprotein [HDL], and low-density lipoprotein [LDL]).

glibenclamide group, 3.2% *Strobilanthes crispus* (SC) leaf extract/ SC1, and 16.8% SC2)

Statistical analysis

Data were performed as mean ± standard deviation (SD) and analyzed by one-way analysis of variance continued with Duncan's multiple range test at $p < 0.05$. The analysis used Statistical Package for the Social Sciences (SPSS) software (version 16.0 SPSS Inc., Chicago, USA).

Results

Feed intake and BW of rat

Feed intake of rats during 14 days of treatment tended to be stable and began to increase at the end period of the study (Figure 1). All treatment groups showed an increase in BW, except PC (Figure 2). This indicated that the treatment of GB, SC1, and SC2 was able to reduce the effect of weight loss experienced by diabetic rats (PC).




Figure 1: Feed intake of rats during 14 days of intervention. Control groups (non-diabetic rat/negative control and streptozotocin [STZ]-induced diabetic rat/positive control), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg body weight of glibenclamide/




Figure 2: Bodyweight (BW) of rats during 14 days of intervention. Control groups (non-diabetic rat/negative control and streptozotocin [STZ]-induced diabetic rat/positive control), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/glibenclamide group, 3.2% *Strobilanthes crispus* (SC) leaf extract/SC1, and 16.8% SC2)

Effect of SC on blood glucose levels

It can be seen from Table 1 that the initial blood glucose level of rats was normal, at 68.4–69.9 mg/dL ($p > 0.05$). Except for NC, almost all treatment groups were given STZ to significantly increase blood glucose levels ($p < 0.05$). After 14 days of treatment, the blood glucose levels of GB, SC1, and SC2 groups were significantly lower than that of day 0 ($p < 0.05$). At the same time, in the PC group, blood glucose levels increased. It was proved that SC has the same effect as the commercially available drug group, and the effective lowering level of blood glucose levels was 3.2% for 14 days of treatment.

Table 1: Effect of various treatments on blood glucose levels of rats

Groups	Blood glucose levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 day)	After intervention (14 days)	
NC	68.5 ± 2.9 ^a	69.8 ± 2.3 ^a	71.3 ± 2.1 [*]	+1.50 ^b
PC	68.7 ± 3.0 ^a	260.0 ± 5.0 ^b	260.9 ± 5.0 [*]	+0.83 ^b
GB	69.9 ± 2.6 ^a	254.7 ± 5.7 ^b	97.0 ± 1.3 [*]	-157.66 ^a
SC1	68.4 ± 2.1 ^a	256.5 ± 5.9 ^b	133.6 ± 3.5 [*]	-122.93 ^a
SC2	69.8 ± 2.6 ^a	257.6 ± 3.6 ^b	109.0 ± 1.4 [*]	-148.6 ^a

*Significantly different between 0 day and 14 days using t-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values shows the decrease and increase of blood glucose levels between 0 day and 14 days of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *Strobilanthes crispus* extract/SC1, and 16.8% SC2), STZ: Streptozotocin, ANOVA: Analysis of variance, NC: Negative control, PC: Positive control, GB: Glibenclamide group, SC: *Strobilanthes crispus*.

Effect of SC on lipid profile

Before STZ administration, the plasma triglyceride levels of rats in all treatment groups were normal, between 74.5 and 76.6 mg/dL. Administering

Table 2: Effect of various treatments on triglyceride levels of rats

Groups	Triglyceride levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 day)	After intervention (14 days)	
NC	76.6 ± 2.5 ^a	78.2 ± 2.0 ^a	81.9 ± 1.9 [*]	+3.74 ^d
PC	75.2 ± 2.7 ^a	126.6 ± 3.6 ^b	127.9 ± 3.6 [*]	+1.33 ^d
GB	74.5 ± 2.0 ^a	127.0 ± 3.6 ^b	97.9 ± 2.5 [*]	-29.23 ^a
SC1	74.6 ± 2.5 ^a	129.4 ± 1.5 ^b	127.6 ± 2.2	-1.86 ^c
SC2	76.0 ± 3.1 ^a	128.4 ± 3.2 ^b	107.1 ± 2.5 [*]	-20.45 ^b

*Significantly different between 0 day and 14 days using t-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values shows the decrease and increase of blood glucose levels between 0 day and 14 days of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *Strobilanthes crispus* extract/SC1, and 16.8% SC2), STZ: Streptozotocin, ANOVA: Analysis of variance, NC: Negative control, PC: Positive control, GB: Glibenclamide group, SC: *Strobilanthes crispus*.

Table 3 shows that the total cholesterol levels in all treatment groups were the same, which were between 94 and 95.4 mg/dL. The presence of STZ induction in the diabetic rat groups significantly increased total cholesterol levels ($p < 0.05$). SC was equivalent to GB in reducing the level of total cholesterol, the best concentration was 16.8% (SC2 group) ($p < 0.05$).

Table 3: Effect of various treatments on total cholesterol levels of rats

Groups	Total cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 days)	After intervention (14 days)	
NC	94.0 ± 2.9 ^a	94.9 ± 2.6 ^a	95.7 ± 2.7	+0.75 ^c
PC	94.4 ± 1.4 ^a	170.5 ± 3.5 ^b	174.4 ± 4.2	+3.86 ^c
GB	95.4 ± 2.2 ^a	177.5 ± 5.1 ^b	128.5 ± 2.1 [*]	-49.07 ^a
SC1	95.2 ± 2.7 ^a	181.0 ± 5.2 ^c	156.8 ± 1.87 [*]	-24.20 ^b
SC2	94.4 ± 3.6 ^a	180.2 ± 7.7 ^c	138.4 ± 2.6 [*]	-41.72 ^a

*Significantly different between 0 days and 14 days using t-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with – and + values shows the decrease and increase of blood glucose levels between 0 days and 14 days of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *Strobilanthes crispus* extract/SC1, and 16.8% SC2), STZ: Streptozotocin, ANOVA: Analysis of variance, NC: Negative control, PC: Positive control, GB: Glibenclamide group, SC: *Strobilanthes crispus*.

LDL cholesterol levels were in the range of 22.4–26.3 mg/dL which belonged to normal category. After being induced, LDL levels in diabetic rats increased and were significantly different compared to the normal rats' group (NC) ($p < 0.05$). The 14-day intervention in the treatment group significantly reduced LDL cholesterol levels ($p < 0.05$), while the untreated diabetes group (PC) did not. SC2 group had the highest

STZ to the diabetic group (PC, GB, SC1, and SC2) had an impact in increasing triglyceride levels in the range of 126.6–129.4 mg/dL. The GB and SC2 interventions were able to significantly reduced blood triglyceride levels with a decrease of 29.23 and 20.45 mg/dL, respectively ($p < 0.05$) (Table 2).

reduction of cholesterol level with the same result with GB in the end of study (Table 4).

Table 4: Effect of various treatments on LDL cholesterol levels of rats

Groups	LDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 days)	After intervention (14 days)	
NC	25.7 ± 1.13 ^a	25.2 ± 1.12 ^a	27.5 ± 1.28 ^a	+6.9 ^d
PC	26.3 ± 1.09 ^b	73.4 ± 2.56 ^c	74.5 ± 1.71	+1.1 ^c
GB	25.2 ± 1.43 ^b	70.2 ± 2.09 ^b	33.4 ± 1.55 ^a	-36.8 ^a
SC1	24.4 ± 1.02 ^a	75.7 ± 1.75 ^c	51.9 ± 1.55 ^a	-23.7 ^b
SC2	22.4 ± 2.47 ^a	75.3 ± 1.46 ^c	38.4 ± 3.11 ^a	-36.9 ^a

*Significantly different between 0 days and 14 days using t-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test. ΔK with - and + values shows the decrease and increase of blood glucose levels between 0 days and 14 days of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *Strobilanthes crispus* extract/SC1, and 16.8% *Strobilanthes crispus* extract/SC2), STZ: Streptozotocin, ANOVA: Analysis of variance, NC: Negative control, PC: Positive control, GB: Glibenclamide group, SC: *Strobilanthes crispus*, LDL: Low-density lipoprotein.

The HDL cholesterol levels of all groups were classified as normal rats at the beginning of the study and ranged from 80.8 to 82 mg/dL. STZ induced a reduction in HDL levels. Treatments for 14 days increased HDL levels significantly ($p < 0.05$) with the same increasing

effect in all groups of rats ($p > 0.05$). Administering SC at a minimum level (3.2%) was able to give the effect of increasing HDL levels which was equivalent to the commercial drug group (GB) (Table 5).

Table 5: Effect of various treatments on HDL cholesterol levels of rats

Groups	HDL cholesterol levels (mg/dL)			ΔK
	Before STZ induction	After STZ induction (0 day)	After intervention (14 days)	
NC	81.9 ± 1.41 ^a	80.1 ± 1.33 ^b	79.5 ± 0.99	+1.10 ^a
PC	81.6 ± 1.07 ^a	25.3 ± 1.43 ^b	25.1 ± 1.20	-0.14 ^b
GB	82.0 ± 1.41 ^a	25.1 ± 0.38 ^b	68.5 ± 1.46 ^a	36.01 ^b
SC1	80.8 ± 2.17 ^a	25.3 ± 0.79 ^b	49.4 ± 1.82 ^a	24.14 ^b

Significantly different between 0 day and 14 days using t-test. Superscript values in the same column are significantly different ($p < 0.05$) using one-way ANOVA followed by multiple comparison Duncan's test.

ΔK with –and+values shows the decrease and increase of blood glucose levels between 0 day and 14 days of intervention, respectively. Control groups (non-diabetic rat/NC and STZ-induced diabetic rat/PC), treatment groups (STZ-induced diabetic rats treated with 5 mg/kg BW of glibenclamide/GB, 3.2% *Strobilanthes crispus* extract/SC1, and 16.8% SC2), STZ: Streptozotocin, ANOVA: Analysis of variance
 NC: Negative control, PC: Positive control, GB: Glibenclamide group, SC: *Strobilanthes crispus*, HDL: High-density lipoprotein.

polyuria (a lot of urination), polydipsia (a lot of drinking), and polyphagia (a lot of eating) with weight loss [19]. SC treatment was able to significantly reduced blood glucose levels in rats and was comparable to the drug (GB group) ($p < 0.05$). These results were in line with the previous study which showed that SC was effective in reducing blood glucose levels by giving the treatment at the dose of 14.7 mg/30 g BW [20]. In another study, processing SC into fermented tea also showed a decrease in blood glucose levels [10]. This could be affected by the flavonoid content in SC which was able to act as an antioxidant by protecting and increasing higher insulin sensitivity [21], [22]. In addition, the antioxidant had the role in accelerating the release of glucose from the circulation which can reduce blood glucose levels [23].

Discussion

Feed intake and bodyweight of rat

In this study, SC treatment did not affect the rats' feed intake, which proved that there was no effect of SC administration on the rats' appetite. During the 14 days of treatment, the weight of rats that received SC intervention also increased a lot, while the weight of untreated diabetic rats (PC) continued to drop. Weight loss in diabetic rats was caused by continual fat catabolism to meet energy needs in their body as a result of the inability of cells to metabolize glucose in the blood. This inability of glucose metabolism could be caused by a disruption of pancreatic cells in producing insulin [15].

Effect of SC on blood glucose levels

The blood glucose levels of rats at the beginning of the study were categorized as normal, then significantly increased to >126 mg/dL or classified as high [16] after STZ induction. STZ induced DM by donating NO which contributed to pancreatic cell damage by increasing the activity of guanylyl cyclase and formation of cGMP. NO was produced when STZ led to metabolism in cells. In addition, STZ was also able to generate reactive oxygen, which had a high role in pancreatic cell damage [17]. In a previous study, the inducement of diabetic by STZ influenced disorientation of liver cellular structures, including glycogen deposition, nuclear location on the periphery of cell membranes, and acidophilic cytoplasm. Liver cell also experienced to **necroticization**, microvascular vacuolization, and hydropic inflammation [18].

In this study, indications of DM in rats could also be observed from the higher urinary frequency (polyuria). In general, people with diabetes have symptoms of

Esterol level which was classified as high because the level was above 10–54 mg/dL [27]. The presence of STZ induction changed the lipid profile to abnormal. As previously discussed, STZ was able to induce an increase in blood glucose levels (Table 1). It could be caused by insulin resistance affecting its fat metabolism [28]. As a result of insulin resistance, sensitive hormone in adipose tissue increased, generated higher lipolysis of triglycerides in adipose tissue, leading to enhancement of free fatty acid levels in the blood. The excess of free fatty acids was converted into phospholipids, cholesterol, and some were used as raw materials for triglycerides generation in the liver which can then be transported into the blood in the form of lipoproteins [29]. In this condition, abnormalities of fat metabolism occurred in the form of increased triglyceride levels, decreased HDL, and increased small dense LDL subtraction, known as the atherogenic lipoprotein phenotype or lipid triad [30]. Therefore, the increase of blood glucose levels in DM patients had a significant correlation with the increase in triglyceride and cholesterol levels.

Administering SC for 14 days had a significant effect in improving lipid profiles. This can be seen from the SC's ability to significantly reduce triglyceride levels at a dose of 16.8% (SC2 group) and total cholesterol or LDL levels at a dose of 3.2% (SC1 group) ($p < 0.05$). However, by administering SC at a dose of 16.8%, the best reduction in total cholesterol and LDL cholesterol levels was achieved or equivalent to commercial drugs. HDL cholesterol levels can be increased to normal and equivalent to commercial drugs starting from the SC dose of 3.2% ($p < 0.05$).

This study was supported by previous experiment that using SC in the form of fermented tea which also showed improvements in lipid profiles such

Proof

AQ3

as lowering cholesterol, triglyceride, and LDL levels, but increasing HDL levels. In the form of SC ethanol extract at a dose of 400 mg/kg BW for 21 days of treatment, there was also a decrease in cholesterol and LDL levels [11]. SC contains bioactive compounds such as alkaloids, saponins, β -sitosterol, flavonoids, tannins, potassium, and polyphenols. β -sitosterol was a hypocholesterolemic substance that might reduce cholesterol levels in the blood [10]. The presence of β -sitosterol and flavonoids controlled cholesterol synthesis by inhibiting the activity of the enzyme acyl-CoA cholesterol acyl transferase in HepG2 cells which play a role in reducing cholesterol esterification in the intestine and liver. It also inhibited the activity of the HMG-CoA enzyme, leading to inhibition of cholesterol synthesis [31]. Flavonoids in SC reduced blood cholesterol levels by reducing or inhibiting the absorption of bile acids and cholesterol in the small intestine. It resulted in high excretion through feces, leading to the formation of bile acids from cholesterol of liver cells. Increased excretion of fecal cholesterol also occurred because of the binding in the intestine between fiber with cholesterol and bile acids which will eventually be excreted or excreted through the feces. This reduced enterohepatic circulation of bile acids and increased the conversion of cholesterol into bile acids, leading to the decrease of plasma cholesterol levels [32].

Conclusions

This study proved that SC extract provided good hypoglycemic effect starting at a concentration of 3.2%. However, administration of 16.8% SC was optimal not only in lowering blood glucose levels but also improving lipid profiles in the form of decreasing levels of triglycerides, total cholesterol, LDL cholesterol, and increasing HDL cholesterol.

Acknowledgments

The authors would like to thank Alma Ata University for the many facilities for this study.

References

- ?
?
?.
2. Yahya J, Stephanie, Wijaya E. Utilization of Keji Beling (*Strobilanthes crispus* BL) Tea as Alternative Medicine of Kidney Stones and Diabetes. Bogor: ???; 2009. 2016;7(4):421-7. <https://doi.org/10.1016/j.jtcme.2016.11.007>
3. Aprilia V, Kurniasari Y, Hadi H, Nurinda E, Ariftiyana S. Effect of Porang Flour (*Amorphophallus oncophyllus*) on Blood Glucose, Triglyceride, HDL, LDL, and Cholesterol of Streptozotocin-induced Diabetic; 2017. AQ4
4. Afifah E. Addition of sapodilla fruit (*Manilkara zapota* L.) decreased the blood glucose levels of diabetic rat (*Rattus norvegicus*). Indonesian J Nutr Diet. 2016;3(3):180.
5. Francisco BM, Salvador M, Amparo N. Oxidative stress in myopia. Oxid Med Cell Longev. 2015;2015:750637. <https://doi.org/10.1155/2015/750637> PMID:25922643
6. Purwaningtyastuti R, Nurwanti E, Huda N. Vitamin C consumption related with blood glucose levels of Type 2-diabetes outpatients. Indonesian J Nutr Diet. 2018;5(1):44.
7. Silalahi M. Utilization of keji beling (*Strobilanthes crispus*) as traditional medicine and its bioactivity. J Edukasi Mat Sains. 2020;9:196-205. AQ5
8. Nurraihana H, Norfarizan-Hanoon NA. Phytochemistry, pharmacology and toxicology properties of *Strobilanthes crispus*. Int Food Res J. 2013;20(5):2045-56.
9. Norfarizan-Hanoon NA, Asmah R, Rokiah MY, Fauziah O, Faridah H. Absence of toxicity of *Strobilanthes crispus* juice in acute oral toxicity study in Sprague Dawley rats. Sains Malaysiana. 2012;41(4):403-9.
10. Fadzelly AB, Asmah R, Fauziah O. Effects of *Strobilanthes crispus* tea aqueous extracts on glucose and lipid profile in normal and streptozotocin-induced hyperglycemic rats. Plant Foods Hum Nutr. 2006;61(1):7-12. <https://doi.org/10.1007/s11130-006-0002-z> PMID:16688478
11. Oktavia S, Eriadi A, Valdis S. Effect of ethanolic extract of keji beling (*Strobilanthes crispus* Blume) leave on cholesterol levels and LDL of hypocholesterolemic male white rats. J Farm Higea. 2018;10(2):110-5.
12. Lim KT, Lim V, Chin JH. Subacute oral toxicity study of ethanolic leaves extracts of *Strobilanthes crispus* in rats. Asian Pac J Trop Biomed. 2012;2(12):948-52. [https://doi.org/10.1016/S2221-1691\(13\)60005-2](https://doi.org/10.1016/S2221-1691(13)60005-2) PMID:23593574
13. Reeves PG, Neilsen FH, Fahey, GC Jr. AIN-93 purified diets for laboratory rodents: Final report of the American institute of nutrition Ad Hoc writing committee on the formulation of the AIN-76A rodent diet. J Nutr. 1993;123(11):1939-51. <https://doi.org/10.1093/jn/123.11.1939> PMID:8229312
14. Ghasemi A, Khalifi S, Jedi S. Streptozotocin-nicotinamide-induced rat model of Type 2 diabetes (review). Acta Physiol Hung. 2014;101(4):408-20. <https://doi.org/10.1556/APhysiol.101.2014.4.2> PMID:25532953
15. Rias YR, Sutikno E. Relationship between body weight and random blood glucose levels of diabetic rats. J Wiyata. 2017;4(1):72-7.
16. Association AD. Classification and diagnosis of diabetes melitus. Diabetes Care. 2015;38 Suppl: S8-16. <https://doi.org/10.2337/dc15-S005> PMID:25537714
17. Nugroho AE. Animal models of diabetes mellitus: Pathology and mechanism of some diabetogenics. Biodivers J Biol Divers. 2006;7(4):378-82.
18. Jaiswal YS, Tatke PA, Gabhe SY, Vaidya AB. Antidiabetic activity of extracts of *Anacardium occidentale* Linn. leaves on n-streptozotocin diabetic rats. J Tradit Complement Med.

- PMid:29034189
19. Snehalata C, Ramachandran A. Diabetes Mellitus in Public Nutrition and Health. Jakarta, Indonesia: Penerbit Buku Kedokteran EGC; 2009.
20. Nonci FY, Leboe DW, Armalia. Activity of ethanolic extract of keji beling (*Strobilanthes crispus* Linn) in decreasing of blood glucose levels of male rats (*Mus musculus*). JF FIK UINAM. 2016;4(1):16-20.
21. Ismawati I, Sembirin LP. The effect of alpha lipoic acid in preventing atherosclerosis lesion on the abdominal aorta of diabetes mellitus Type 2 rats. J Kedokt dan Kesehat. 2020;16(1):???
22. Ashari A, Nurinda E, Fatmawati A. Increase of insulin levels of Streptozotocin-induced male Wistar rats (*Rattus norvegicus*) after ethanolic extract treatment of brotowali stem (*Tinospora crispa* L.). Indonesian Pharm Nat Med J. 2021;5(1):1.
23. Widowati W. Potency of antioxidant as antidiabetes. J Kesehat Masy. 2008;7(2):193-202.
24. Suryaatmadja M. Triglyceride test without fasting. Summit Lipid Update. 2012;7:1-2.
25. Nugraheni M, Lastariwati B, Purwanti S. Proximate and chemical analysis of gluten-free enriched, resistant starch Type 3 from Maranta arundinacea flour and its potential as a functional food. Pak J Nutr. 2017;16(5):322-30.
26. Pradana DA, Dwiratna DW, Widyarini S. Activity of standardized-ethanolic extract of red spinach (*Amaranthus tricolor* L.) as preventive efforts of steatosis: *In vivo* study. J Sains Farm Klin. 2017;3(2):120.
27. Nurhidajah, Astuti R, Nurrahman. Black rice potential in HDL and LDL profile in Sprague Dawley rat with high cholesterol diet. IOP Conf Ser Earth Environ Sci. 2019;292(1):012019.
28. Trivedi NA, Mazumdar B, Bhatt JD, Hemavathi KG. Effect of shilajit on blood glucose and lipid profile in alloxan- induced diabetic rats. Indian J Pharmacol. 2004;36(6):373-6.
29. Yassin MM, Mwafy SN. Protective potential of glimepiride and Nerium oleander extract on lipid profile, body growth rate, and renal function in streptozotocin-induced diabetic rats. Turkish J Biol. 2007;31(2):95-102.
30. Adam J. Dislipidemia: Internal Medicine Science. 4th ed. Jakarta, Indonesia: Pusat Department of Internal Medicine Science, Medicine Faculty. Universitas Indonesia; 2006.
31. Riyanto S, Muwarni RH. Black soyghurt lowered LDL levels of hypercholesterolemic rats. Indonesian J Nutr Diet. 2016;3(1):1.
32. Potter SM. Overview of Proposed Mechanisms for the hypocholesterolemic effects os soy. J Nutr. 1995;125:606.

Author Queries???

AQ1: Kindly provide history details

AQ2: Please check the term.

AQ3: Kindly provide page number AQ4:

Kindly provide publisher details AQ5:

Kindly provide in English language