



Plagiarism Checker X Originality Report

Similarity Found: 9%

Date: Friday, December 15, 2023

Statistics: 497 words Plagiarized / 5265 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

Full Terms & Conditions of access and use can be found at

<https://www.tandfonline.com/action/journalInformation?journalCode=iijf20>

International Journal of Food Sciences and Nutrition ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/iijf20> Intra-uterine exposure to lower maternal haemoglobin concentration is associated with impaired cognitive function in stunted adolescents: results from a 17-year longitudinal cohort study in Indonesia Muhammad Asrullah, Hastrin Hositanisita, Monique L'Hoir, Siti Muslimatun, Edith JM Feskens & Alida Melse-Boonstra To cite this article: Muhammad Asrullah, Hastrin Hositanisita, Monique L'Hoir, Siti Muslimatun, Edith JM Feskens & Alida Melse-Boonstra (2023) Intra-uterine exposure to lower maternal haemoglobin concentration is associated with impaired cognitive function in stunted adolescents: results from a 17-year longitudinal cohort study in Indonesia, International Journal of Food Sciences and Nutrition, 74:8, 826-835, DOI: 10.1080/09637486.2023.2267792 To link to this article: <https://doi.org/10.1080/09637486.2023.2267792> © 2023 The Author(s). Published with license by Taylor & Francis Group, LLC. View supplementary material Published online: 12 Dec 2023.

Submit your article to this journal Article views: 531 View related articles View Crossmark data STUDIES IN HUMANS International Journal of food ScienceS and nutrItIon 2023, Vol. 74, no . 8, 826–835 Intra-uterine exposure to lower maternal haemoglobin concentration is associated with impaired cognitive function in stunted adolescents: results from a 17-year longitudinal cohort study in Indonesia Muhammad Asrullah a,b, Hastrin Hositanisita a,c, Monique L'Hoir a, Siti Muslimatun d, Edith JM Feskens a and Alida Melse-Boonstra a a division of Human nutrition and Health, Wageningen university and r esearch, the netherlands; bcentre for Health Policy and Management, faculty of Medicine Public Health and nursing, universitas Gadjah Mada, Yogyakarta, Indonesia;

Department of Nutrition, Faculty of Health Sciences, the University of Airlangga, Yogyakarta, Indonesia; Department of Food Science and Nutrition, Indonesia International Institute for Life Sciences, Jakarta, Indonesia

ABSTRACT The long-term impact of maternal anaemia on cognitive performance remains unknown.

Indonesian longitudinal cohort data of 363 paired pregnant mothers and their 10–14-year-old offspring were used to investigate the association between maternal haemoglobin (Hb) concentration and their offspring's cognitive function (assessed by Raven's Progressive Matrices test) during adolescence. The weighted anaemia prevalence was 49.3% in pregnant mothers and 22.2% in adolescents. Adolescents who were stunted, anaemic, or living in a rural area had significantly lower cognitive scores than their counterparts. Maternal Hb was not associated with adolescent cognitive function (β : 0.14; 95%CI: -0.052–0.340).

However, the effect of maternal Hb concentration on offspring's cognitive function was modified by stunting status (β , stunted: 0.44; 95%CI: 0.05–0.82; non-stunted: 0.01; 95%CI: -0.02–0.24). This study shows adverse cognitive outcomes at adolescent age are likely multi-causal and can be partially explained by intra-uterine exposure to low maternal Hb concentrations. Introduction About one-third of the world's population, especially in Low-and Middle-Income Countries (LMICs), is suffering from iron deficiency anaemia, a condition with abnormally small-sized red blood cells and reduced haemoglobin (Hb) concentration (Chaparro and Suchdev 2019). Anaemia affects around 800 million children and women in the world, with almost half of all pregnant women being affected (World Health Organization 2011).

In Indonesia, anaemia is considered a moderate public health problem with a prevalence among pregnant women of 37.1% in 2013, which increased to 48.9% in 2018 based on the Indonesian National Basic Health Survey (Kementerian Kesehatan Republik Indonesia 2018). Pregnancy is a vital stage in human development, and maternal anaemia during the intra-uterine period can affect growth and development of the offspring.

A meta-analysis of 26 cohort studies (both with prospective and retrospective study designs) conducted in LMICs confirmed that anaemia during pregnancy is associated with low birth weight, preterm birth, and perinatal mortality (Rahman et al. 2016). A cohort study in India among 1,007 pregnant women found that women who were anaemic in the second trimester of gestation gave birth to infants with lower birth weight (Nair et al. 2016). Low birth weight is an important determinant of stunting later in life, as also shown for children aged 0–23 months in Indonesia based on three waves of data (2000, 2007, and 2014) from the Indonesian Family Life Survey (IFLS) (Hanifah

et al. 2018).

Previous studies also found that maternal anaemia is associated with anaemia in the offspring. For instance, a study in China found that © 2023 the author(s). Published with license by Taylor & Francis Group, LLC. CONTACT Muhammad Asrullah Muhammad.asrullah@wur.nl division of Human nutrition and Health, Wageningen university and research, Stippeneng 2, 6708 WE Wageningen, the Netherlands. Supplemental data for this article can be accessed online at <https://doi.org/10.1080/09637486.2023.2267792>. <https://doi.org/10.1080/09637486.2023.2267792> this is an open access article distributed under the terms of the creative commons attribution license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

the terms on which this article has been published allow the posting of the accepted Manuscript in a repository by the author(s) or with their consent. **ARTICLE HISTORY** Received 4 July 2023 Revised 30 August 2023 Accepted 2 October 2023 **KEYWORDS** Pregnancy; anaemia; adolescent; cognition; Indonesia **INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION** 827 infants at the age of 5–7 months and 11–13 months had a higher risk of anaemia when their mothers had been anaemic during mid-pregnancy (Chang et al. 2013).

An intervention study among 359 Filipino pregnant women found that maternal iron deficiency anaemia is linked to an increased risk of iron deficiency during infancy (Mendez and Adair 1999). More importantly, children tend to have delayed neurocognitive development, including mental and motor development, at young age if their mothers had been anaemic during pregnancy (Tran et al. 2014). This may be due to the non-reversible nature of neurodevelopmental damage incurred during early stages of life (Beard 2008). Also, anaemia during infancy, which is predominantly caused by iron deficiency worldwide (De Benoist et al.

2008), is associated with poorer cognitive and motor development (Grantham-McGregor and Ani 2001; Sania et al. 2019). The effect of low haemoglobin on cognitive function may occur in intra-uterine of the mother during pregnancy. The Intra-uterine maternal anaemia can be a predisposing risk factor for intra-uterine infant growth such as brain development and cognitive function. It is because insufficient levels of haemoglobin hinder the proper circulation of oxygen within the body and leads to a state of oxidative stress or persistent hypoxia, creating conditions that may contribute to limitations in foetal growth (Kozuki et al. 2012). The potential mech -

anism underlying an iron deficiency-related cognitive deficit is not yet well studied in humans (Veena et al.

2016), but animal studies showed that iron deficiency may interfere with neurogenesis, myelination, dendrite growth, and protein expression during neurodevelopment (Beard et al. 2006; Lozoff et al. 2006; Fretham et al. 2011). A study in India showed that 3-week old infants whose mothers had not been anaemic during pregnancy scored 3.9 times higher on behavioural outcomes such as attention and interactive abilities (Menon et al. 2016). A nationwide prospective cohort study over 30 years conducted among 11,656 Finnish women found that low maternal Hb, especially during late gestation, was associated with lower educational performance of their offspring at the ages of 14 and 16 years (Fararouei et al. 2010). Adolescence is a period of distinctive neurocognitive growth in which a basic reorganisation of the brain occurs (Giedd et al. 1999).

Evidence on the persistent impact of early-life anaemia on cognitive performance in later life has been summarised in a systematic review including 27 observational and randomised controlled trials (RCT), but it was concluded that there is a scarcity of studies with long-term follow-up into adolescence and adulthood (Alwan and Hamamy 2015; Janbek et al. 2019). This piece of information can help to even better understand the long-term intergenerational consequences of maternal anaemia. We therefore aimed to evaluate the relationship between intra-uterine maternal Hb concentration and cognitive function of their offspring at adolescent age (10–14 years), using longitudinal data from the Indonesian Family Life Survey (IFLS).

Methods Indonesian Family Life Survey (IFLS) IFLS is a longitudinal survey in Indonesia that started in 1993/1994 (IFLS 1). The IFLS used a multistage stratified sampling design and collected data at individual and household level (Strauss et al. 2016). The total sample of IFLS comprises >30,000 individuals from 13 out of 27 provinces in Indonesia and represents about 83% of the Indonesian population. IFLS 2 was performed in 1997/1998, IFLS 3 in 2000, IFLS 4 in 2007/2008, and IFLS 5 in 2014/2015. For the current analysis, we only used data from IFLS waves 2 to 5, since maternal Hb concentration was not assessed in the first wave.

Participants Subjects were adolescents aged 10–14 years from IFLS waves 4 and 5, who had a complete cognitive score record ($n = 8,109$). Participants without weight and height measurements ($n = 22$) and Hb data ($n = 349$) were excluded, yielding 7,712 adolescents. A total of 867 pregnant women >15 years with complete Hb data from waves 2 and 3 were then paired with their children at adolescent age ($n = 363$; Figure 1).

Assessment of cognitive function Cognitive function was measured using an age-appropriate subset of questions from Raven's Progressive Matrices (RPM). RPM is the most commonly used nonverbal test to measure general intellectual ability, or cognitive function (Sandjaja et al. 2013). It relates to several abilities such as analysing, reasoning, and understanding of unconcreted concepts (Prabhakaran et al. 1997).

This intelligence test has been validated for use in different socio-economic groups, ethnicities, and cultures (Raven 2000). Two members of the testing division of the Indonesian Ministry of Education adapted the test for the Indonesian setting by drawing items from the National Achievement Test (Frankenberg and Thomas 2000). In IFLS, an RPM module (EK1) was developed for 828 M. ASRULLAH ET AL. participants aged 7–14 years while a larger module including EK1 items was developed for participants aged 15 years or older.

The subset EK1, as used for the present study, consists of 17 questions; 12 questions to test cognitive abilities, such as general intelligence and non-concrete reasoning, and 5 questions to test mathematic skills. Correct answers were scored as "1" and incorrect answers as "0", so that the maximum total score was 17 (Supplementary Table 1). Assessment of maternal and adolescent anaemia Hb concentrations were determined from finger prick blood by trained health workers using a point-of-care Hb test (HemoCue Hb201+, Ängelholm, Sweden), as described before for IFLS (Strauss et al. 2016).

For adolescents, Hb was measured when they were 10–14 years old. Based on WHO guidance, anaemia status was defined as Hb <11.5 g/dl for adolescents aged 10–11 years, < 12.0 g/dL for adolescents aged 12–14 years, and <11 g/dL for pregnant women (World Health Organization 2013). In the present study, Hb concentrations were not adjusted for altitude or smoking due to the unavailability of such data. Assessment of other variables Body Mass Index (BMI)-for-age and height-for-age z-scores were calculated using the WHO growth standard for children aged 5–19 years, with a BMI-for-age z-score <-2 SD classified as thin, =-2 SD to =1 SD as normal body weight, >1 SD as overweight, and >2SD as obese. Height-for-age z score (HAZ) <-2 SD defined stunting (World Health Organization 2007). Per capita expenditure (PCE) was used to indicate socioeconomic status (SES) (Preedy 2012).

The PCE was formulated based on all household expenditures (either for food and non-food items, education and housing) over a month, divided by the number of household members (Witoelar 2009). To correct for inflation rate and to standardise the value, PCE of waves 2 and 3 were converted to Jakarta's price level in December 2000

(Witoelar 2009). PCE missing values ($n = 2$) were imputed with the median value of PCE in the sub-district where they lived. PCE is presented as tertiles, with the first tertiles representing the lowest per capita expenditure class (Vaezghasemi et al. 2016).

Ethical approval The IFLS and the survey protocol were reviewed and approved by Institutional Review Boards (IRB) in the USA and the Ethics committee of Universitas Indonesia based in Jakarta for the first wave of IFLS, and by Universitas Gadjah Mada in Yogyakarta for IFLS waves two to five. The ethical clearance number from RAND's Human Subjects Protection Committee (RAND's IRB) was s0064-06-01-CR01. Figure 1. flowchart of participant selection based on data availability. **INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION** 829 All methods were performed in accordance with the relevant guidelines and regulations by including a statement in the Ethics approval and informed consent paper.

Informed consent was obtained from all participants to participate after explanation of the study. **Statistical analysis** The statistical analysis was performed using STATA version 16 (StataCorp. 2019). Data analysis was done based on complete cases (Figure 1). A comparison was made between included ($n = 363$) and excluded cases ($n = 9$) to verify that this approach did not lead to selection bias (Supplementary Table 2). Subjects' characteristics such as age, sex, living area, BMI z-score, and HAZ were first analysed descriptively.

Before regression analysis, the assumption of normal value distribution was checked for continuous data by inspecting the Q-Q plot of regression standardised residuals, and multicollinearity was checked by variance inflation factor ($VIF < 10$), with $VIF < 10$ reflecting no collinearity. A one-way ANOVA was performed to analyse differences in mean total cognitive scores between groups based on household variables (SES and living area), maternal variables (age and anaemia status), and adolescent characteristics (sex and BMI-for-age z-score). Multiple linear regression was used to relate maternal Hb concentration during pregnancy to adolescents' total cognitive score before and after adjusting for maternal age, maternal height, and SES.

The adjustment was based on a Directed Acyclic Graph (DAG) in which maternal age, maternal height, and SES were found to be confounders of the direct effect of maternal anaemia on adolescent cognitive function (Supplementary Figure 1). Adolescent characteristics, such as Hb concentration, HAZ, and BMI-for-age-z-score were tested to be effect modifiers or mediating factors, since these factors may either be more proximate predictors in the pathway between maternal Hb and cognitive outcomes, or they may amplify or downplay any effects.

This was done by including interaction terms between maternal Hb concentration and children's Hb concentration, HAZ, and BMI-for-age-z-scores to identify the effect modifiers, and a mediation analysis (Mehmetoglu 2018) to identify the mediating factors. Variables with a p-value <0.05 in the interaction model were used for stratification to estimate the group-specific regression coefficients with their 95% confidence intervals. Results In Table 1 , the characteristics of the 363 paired mothers-adolescents are summarised. Almost half of the pregnant women (49.3%) were anaemic. Anaemia prevalence among adolescents was 22.2%, and 32.3% of the adolescents were stunted.

The mean cognitive score of the adolescents was 12.35 out of 17 (SD: 2.95). Adolescents who were stunted, anaemic or lived in a rural area had lower cognitive test scores than their counterparts who were non-stunted (11.68 ± 3.01 vs 12.66 ± 2.86 , $p = 0.003$), non-anaemic (11.33 ± 3.28 vs 12.66 ± 2.74 , $p < 0.001$), or lived in an urban area (11.92 ± 3.27 vs 12.68 ± 2.62 , $p = 0.015$). Cognitive score did not differ between adolescents whose mothers had been anaemic or non-anaemic during pregnancy, nor for any of the other characteristics (Table 2).

No association was found between maternal Hb and adolescent's cognitive score in unadjusted and adjusted models (Table 3). We did not find any mediation of adolescent Hb, BMI-Z score, and HAZ in the association between maternal Hb and adolescent cognitive scores (Supplementary Table 3). However, we observed an interaction between maternal haemoglobin concentration and stunting ($p = 0.039$), as well as a difference in cognitive scores between stunted and non-stunted adolescents among the group of anaemic Table 1.

characteristics of pregnant women and their 10–14 y old adolescents ($n = 363$) from IfIS waves 2–5 who were included in the data analysis. characteristics Valuesa Household Per capita expenditure, Idr low 91 (25.07) Medium 182 (50.14) High 90 (24.79) living area urban 214 (58.63) rural 148 (40.55) Pregnant women age, y <20 y 1. (13.77) 20–35 y 279 (76.86) >35y 33 (9.09) Haemoglobin concentration, g/dl 11.06 ± 1.55 a naemic 179 (49.31) adolescents age, y 13.17 ± 1.24 Sex (boys/girls) 179 (49.31)/184 (50.69) BMI-for-age z-score 0.002 ± 1.00 t hin 0 (0) normal weight 316 (87.05) o verweight 28 (7.71) obese 19 (5.23) Height-for-age z-score -1.52 ± 1.02 Stunted 117 (32.23) Haemoglobin concentration, g/dl 12.90 ± 1.47 a naemic 81 (22.19) aValues are means $\pm$ Sd for continuous variables, and n (%) for categorical data; IfIS: Indonesian family life Survey; Idr: Indonesian rupiah based on d ecember 2000 Jakarta baseline. 830 M. ASRULLAH ET AL. mothers (Supplementary Table 4).

Therefore, we continued with stratifying the analysis for stunted and non-stunted

adolescents separately. In contrast to the non-stunted population, we found a positive association - **tion between maternal haemoglobin concentration** and cognitive score in stunted adolescents both in the unadjusted model ($\beta = 0.46$, 95%CI 0.08–0.84) as well as in the model adjusted for maternal age, maternal height, and socioeconomic status ($\beta = 0.44$, 95%CI 0.05–0.82) (Table 3). No interaction was observed for maternal Hb and adolescent BMI ($p = 0.830$), and maternal Hb and adolescent Hb ($p = 0.844$).

Discussion **In this study, we examined the association between maternal Hb** concentration and her offspring's cognitive function presented as RPM total scores at adolescent age (10–14 years).

We found that adolescents who suffered from anaemia were stunted or lived in a rural area had lower total cognitive scores. Maternal Hb **was not associated with** adolescent cognitive function. However, **the association between maternal Hb** concentration and cognitive function of their offspring was modified by stunting status of the adolescents and showed a positive association in stunted adolescents indicating synergy. **Our study showed that** almost half of the pregnant women were anaemic (49.3%), and the overall anaemia prevalence in adolescents was 22.9%.

This shows that, based on WHO classification, anaemia was a moderate (adolescents) to severe (pregnant women) public health problem in the population under study (De Benoist et al. 2008). **It should be noted that** the data from pregnant women in this study originated from IFLS waves 2 and 3, conducted in 1997 and 2000. Another study also reported **that the prevalence of** anaemia during pregnancy in this period was high (45.1–46.5%) (Barkley et al. 2015). An explanation for the high prevalence of maternal anaemia during this time period could be the Indonesian economic crisis, in which the poverty increased from 17% to 24% from 1996 to 1999 (Rizky et al. 2019).

Although there has been a dip in the prevalence of maternal anaemia recorded by **the national basic health survey** (Riskesdas; Riset Kesehatan Dasar) in 2013, i.e. 37.1%, data from the most recent Riskesdas conducted in 2018, shows a similarly high prevalence of maternal anaemia, i.e. 48.9%, despite the fact that a nationwide iron-folic acid (IFA) supplementation program has been implemented since 1990 (Kementerian Kesehatan Republik Indonesia 2018).

Incomplete coverage of the IFA supplementation program, with a national coverage of ~ 78% from 2002 to 2007, might be a reason for the lack of decline in the prevalence of maternal anaemia. It should also be noted that IFA Table 2. Mean ($\pm$ Sd) total cognitive score on the Raven's **Progressive Matrices (rPM) test** of adolescents ($n = 363$) from IFLS in relation to household, maternal and individual characteristics. characteristics cognitive score (mean $\pm$ Sd) p Value Household Per capita expenditure, Idr low 12.07 $\pm$

2.89 0.581 Medium 12.43 ± 3.01 High 12.46 ± 2.89 living area 0.015 urban 12.68 ± 2.62 rural 11.92 ± 3.27 Pregnant women age, y 11.71 ± 2.84 0.171 20–35 y 12.50 ± 2.92 12.03 ± 3.26 Haemoglobin concentration a naemic 12.15 ± 2.88 0.204 non-anaemic 12.66 ± 2.74 adolescents Sex Boys 12.47 ± 2.93 0.426 Girls 12.22 ± 2.97 BMI-for-age z-score t hin – 0.212 normal weight 12.26 ± 2.99 o verweight 13.28 ± 2.62 obese 12.31 ± 2.45 Height-for-age z-score Stunted 11.68 ± 3.01 0.003 non-stunted 12.66 ± 2.86 Haemoglobin concentration, g/dl a naemic 11.33 ± 3.28 0.000 non-anaemic 12.66 ± 2.74 IfIS: Indonesian family life Survey; Idr: Indonesian rupiah currency based on d ecember 2000 Jakarta baseline. Table 3.

Multiple linear regression of **the association between maternal haemoglobin concentration** (g/l) during pregnancy and cognitive score of her adolescent child aged 10–14 years from IfIS waves 2–5. cognitive score a unadjusted adjustedb β (95% cI) p-Value β (95% cI) p-Value total study population (n = 363) 0.14 -0.06, 0.34 0.165 0.14 -0.05, 0.34 0.160 Stunted adolescents (n = 117) 0.46 0.08, 0.84 0.018 0.44 0.05, 0.82 0.026 non-stunted adolescents (n = 246) 0.01 -0.22, 0.23 0.960 0.01 -0.22, 0.24 0.922 acognitive score was assessed using the eK1 module from the raven's Progressive Matrices test in IfIS.

bModel was adjusted for maternal age, maternal height, and socioeconomic status. **INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION** 831 supplementation has not been distributed equally over the population, i.e. pregnant women who were higher educated and those who lived in urban areas took IFA supplementation more often than pregnant women with a lower educational level and who lived in rural areas (Fiedler et al. 2014). Populations **living in rural areas** often experience more health disparities (Irianti and Prasetyoputra 2021). This may be related to geographic isolation, lower SES, limited access to medical and preventive care, and less job opportuni - ties.

A previous study confirmed that low SES was associated with anaemia during pregnancy in Indonesia (Chandra et al. 2018). Our finding on the prevalence of anaemia among adolescents is largely in line with the most recent Riskesdas (2018), showing **that the prevalence of** anaemia was 26.8% among those aged 5–14 years and 32.4% among those aged 15–24 years (Kementerian Kesehatan Republik Indonesia 2018).

Anaemia among adolescents can be due to the increased need for iron and other micronutrients in support of their physical growth and, for girls, to compensate for menstrual loss of iron (de Benoist et al. 2008) and the adoption **of unhealthy eating habits** in a time period of physical, mental, and emotional change (Agustina et al. 2020). We did not find an overall **association between maternal Hb** concentration and total cognitive scores among adolescents either in unadjusted or adjusted models.

The limited number of studies on this topic performed to date have shown inconsistent findings (Alwan and Hamamy 2015; Janbek et al. 2019). Some observational studies found a negative/inverse association (Hernández-Martínez et al. 2011; Tran et al. 2014; Aranda et al. 2017), or an inverted U-shaped association (Mireku et al. 2015) when developmental outcomes were measured in infancy. The largest 30-year prospective cohort study to date, including 11,656 Finnish women, showed that low maternal haemoglobin, especially during late pregnancy, was linked to lower educational achievement of their offspring at the age of 14 and 16 years (Fararouei et al. 2010). This is in line with study findings from 850 women and their children in China, showing that maternal anaemia, especially in the third trimester, was associated with worse neurodevelopment in their children (Chang et al. 2013).

We could not differentiate between anaemia across trimesters of pregnancy since data on gestational age at the time point of Hb testing were not collected. Discrepancies between study findings may also be explained by certain contextual differences, such as differences in SES, prevalence of malnutrition, health status, and health care access. It should be emphasised that Hb cannot be interpreted as a direct measure of iron deficiency, and therefore it is also worthwhile to take a look at studies that used indicators of iron status.

Iron plays an important role during foetal development in cell maturation and myelination of the frontal lobes (Todorich et al. 2009). Iron-containing enzymes are crucial especially in the development of the striatum and hippocampus, (Alwan and Hamamy 2015) which are important for memory, optimal learning, processing speed, and decision making (O'Shea et al. 2016). Iron deficiency during specific time windows of pregnancy may importantly modify in utero neurodevelopment (Alwan and Hamamy 2015).

A rapid growth of the hippocampus in humans starts in late pregnancy throughout the first year of life (Alwan and Hamamy 2015). There is some evidence that low maternal iron status especially in the late gestational period is adversely associated with offspring's neurodevelopment from infancy (<1 year) till adulthood (>18 years) (Menon et al. 2016). Furthermore, iron deficiency together with low haemoglobin concentration is more clearly associated with long-term deficits in cognitive and motor function of children than low iron status only.

Children with iron deficiency and haemoglobin concentration = 105 g/L had consistently lower cognitive abilities as compared to non-anaemic iron deficient children (Carter et al. 2010). Moreover, in children with iron deficiency and low haemoglobin

concentration (Hb < 115 g/L for 5–11 years old; Hb < 120 g/L for 12–13 years old), still a positive association between haemoglobin concentration and cognitive performance was shown, although no such association existed in non-anaemic iron-deficient children (Sungthong et al. 2002). An RCT found a beneficial effect of correction of iron status during late pregnancy on developmental child outcomes (Chang et al. 2013).

In the present study, we did not have information on iron metabolism indicators and therefore could not assess the association between maternal iron status and cognitive outcomes. The present study only showed a significant association between maternal Hb and cognitive function for adolescents who were stunted. Collective data from 137 countries found that maternal anaemia is one of the key factors of child stunting (Danaei et al. 2016).

About one-third of adolescents in our study were stunted, and it has been shown before that stunted children also have a higher risk of being anaemic (Rahman et al. 2019; Wang et al. 2020). Moreover, stunting has been associated with a lower cognitive functioning at school age and adulthood, along with other risk factors, such as poverty and 832 M. ASRULLAH ET AL. poor home environment (Black et al. 2017). A prospective birth cohort study among 1,291 children conducted in eight countries in Asia, America, and Africa showed that early-onset persistent stunting was associated with lower cognitive development scores at 5 years of age (Alam et al.

2020). The effects might persist to adolescent age since a study from the Philippines showed lower cognition scores at early adolescent age (8 and 11 years) after recovering from stunting (Mendez and Adair 1999). Another study found that stunted Jamaican children still had lower cognitive scores at adolescent age (17–18 years) despite nutritional supplementation and stimulation at the age of 9–24 months (Walker et al. 2005). Thus, stunting may be an important explanatory factor in the pathway between maternal anaemia and cognitive function.

Nevertheless, our analysis revealed that stunting was rather an effect modifier than a mediating factor between maternal Hb and cognitive outcomes at adolescent age. The results suggested that stunting exacerbates the effect of maternal anaemia on adolescent cognitive function. Since adolescents in the present study were born during the economic crisis, their parents' economic status may have been affected. The Young Lives study, an observational cohort study among 3000 children in four developing countries (Ethiopia, Peru, India, and Vietnam), found that three factors were strongly associated with cognitive function at adolescent age: child growth (including stunting status), parental education and socioeconomic status (Crookston et al. 2014).

Previous studies in Indonesia found a stronger and more consistent association of adolescents' cognitive scores with socioenvironmental risk factors including stunting than with biomedical factors (Prado et al. 2017). Previous IFLS studies found only a weak association of household expenditure with children's cognitive function (Maika et al. 2015; Sartika et al. 2019). It should be noted that stunting in adolescents is multi-causal and can be the result of infection, poor nutrition, and environmental stress starting from the foetal period to later in life (Christian and Smith 2018).

Furthermore, inequalities in education and learning outcomes exist, with rural areas lagging behind compared to urban areas in Indonesia (The World Bank 2017). Thus, the lower cognitive function of adolescents who were both exposed to maternal anaemia and were stunted in this study may be attributable to the same root causes of long-term poverty. To our knowledge, this is one of few studies which evaluated the long-term effects of maternal anaemia on cognitive function in adolescence, using a prospective longitudinal study design.

However, gestational age at the time point of Hb testing was not assessed, and no indicators of iron status were measured in this study. Furthermore, this study also did not take into account the altitude and smoking status to correct Hb concentrations as suggested by the WHO (World Health Organization 2013) due to unavailability of such data. There may also be unmeasured confounding, e.g. by other maternal exposures and variables that impact both maternal nutritional status and child development.

In that case, there is a possibility that maternal anaemia may not be causal to poor cognitive development of her stunted adolescent child, but rather is a result of the same cause, i.e. poverty. The RPM is a standardised method to measure non-verbal cognitive ability that has previously been used in Indonesia (Sandjaja et al. 2013) and is not biased by educational background or by cultural or linguistic deficiencies (Motta and Joseph 2000), but we acknowledge that we cannot compare cognitive function measured by RPM with other tools.

In conclusion, maternal Hb concentration was, overall, not associated with cognitive function in adolescents aged 10–14 years, but it was associated in those who were stunted. For each 1 g/dL increase in Hb, cognitive scores increased by 0.44 in stunted adolescents, but not in non-stunted ones. This study reveals that adverse cognitive outcomes at adolescent age is likely to be multi-causal. Authors' contributions All authors had an essential role in formulation of the research questions; MA, HH, and AMB wrote the first draft of the paper, MA analysed the data; MA, MLH, SM, AMB, and EJM were involved in interpretation of the data and revision of the manuscript. All authors have read and approved the final paper. Disclosure statement No potential

conflicts of interest are reported by the author(s).

Funding This study was performed as part of the PhD studies of the first author (MA) for which a scholarship from Lembaga Pengelola Dana Pendidikan (LPDP), Republic of Indonesia was received. LPDP had no role in the design, data analysis, or interpretation of the data and writing process of this article. INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION 833 Data availability statement The data are available online at <https://www.rand.org/>

INTERNET SOURCES:

1% - <https://www.tandfonline.com/toc/ijf20/74/8>
<1% - <https://www.tandfonline.com/journals/ijf19>
<1% - <https://europepmc.org/article/MED/35361192>
<1% - <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0288073>
<1% - https://faseb.onlinelibrary.wiley.com/doi/abs/10.1096/fasebj.29.1_supplement.729.21
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9501636/>
<1% - <https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-021-11014-1>
<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9559178/>
<1% - https://handbook-5-1.cochrane.org/chapter_19/19_1_1_what_is_a_prospective_meta_analysis.htm
<1% - <https://europepmc.org/article/MED/28588921>
<1% - <https://www.who.int/data/nutrition/nlis/info/low-birth-weight>
<1% - <https://e-journal.unair.ac.id/MGI/search/authors/view?givenName=Sandy&familyName=Ardiansyah&affiliation=Division%20of%20Human%20Nutrition%2C%20Wageningen%20University%20and%20Research%2C%20Wageningen%2C%20Netherland&country=&authorName=Ardiansyah%2C%20Sandy>
<1% - <https://www.cambridge.org/core/services/aop-cambridge-core/content/view/D19106D27A66224BEC314B293F813BAF/S2513843X23000270a.pdf/maternal-grandmothers-buffer-the-effects-of-ethnic-discrimination-among-pregnant-latina-mothers.pdf>
<1% - <https://www.jstor.org/stable/26581916>
1% - <https://www.tandfonline.com/doi/pdf/10.1080/19475683.2023.2226196>
<1% - <https://www.tandfonline.com/doi/pdf/10.1080/21645515.2023.2257048?needAccess=true>

<1% - <https://www.nature.com/articles/s41372-019-0375-0>
 <1% - <https://pubmed.ncbi.nlm.nih.gov/34394714/>
 <1% - <https://www.sciencedirect.com/science/article/pii/S0022316622146730>
 <1% - <https://pubmed.ncbi.nlm.nih.gov/31062441/>
 <1% - <https://www.nejm.org/doi/full/10.1056/NEJMoa1700732>
 <1% -
https://www.researchgate.net/publication/256478548_Is_Adolescence_a_Sensitive_Period_for_Sociocultural_Processing
 <1% - <https://www.sciencedirect.com/science/article/pii/S089085672100366X>
 <1% - <https://dornsife.usc.edu/johnstrauss/ifls/>
 <1% -
https://www.rand.org/content/dam/rand/pubs/working_papers/WR1100/WR1143z2/RAND_WR1143z2.pdf
 <1% -
[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)01259-9/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)01259-9/fulltext)
 <1% - <https://www.frontiersin.org/articles/10.3389/fnint.2021.719076/full>
 <1% -
<https://opengovasia.com/indonesia-government-to-launch-programmes-to-enhance-the-education-sector/>
 <1% - <https://bestpractice.bmj.com/topics/en-gb/93>
 <1% - <https://www.e-cep.org/upload/pdf/cep-2020-01508.pdf>
 <1% - <https://www.nature.com/articles/s41598-021-02464-z>
 <1% -
<https://archpublichealth.biomedcentral.com/articles/10.1186/s13690-023-01138-8>
 <1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9097111/>
 <1% -
https://www.researchgate.net/publication/359878132_Secular_trends_in_Japanese_adult_height_the_roles_of_environment_and_educational_attainment
 <1% -
https://en.wikipedia.org/wiki/International_Journal_of_Food_Sciences_and_Nutrition
 <1% - <https://www.springer.com/gp/editorial-policies/informed-consent>
 <1% -
<https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-020-10001-2>
 <1% -
https://www.researchgate.net/publication/335626297_Principle_Assumptions_of_Regression_Analysis_Testing_Techniques_and_Statistical_Reporting_of_Imperfect_Data_Sets
 <1% - <https://openstax.org/books/statistics/pages/13-1-one-way-anova>
 <1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8974490/>
 <1% - [https://jn.nutrition.org/article/S0022-3166\(23\)35483-X/fulltext](https://jn.nutrition.org/article/S0022-3166(23)35483-X/fulltext)
 <1% - <https://academic.oup.com/ejcts/article/55/2/179/5265263>

<1% -

https://www.researchgate.net/publication/367259368_Gestational_age_at_birth_and_cognitive_outcomes_in_adolescence_population_based_full_sibling_cohort_study

<1% - <https://bmcpsy psychiatry.biomedcentral.com/articles/10.1186/s12888-019-2082-y>

<1% - <https://www.sciencedirect.com/science/article/pii/S266614972300021X>

<1% - <https://www.mdpi.com/2072-6643/15/15/3420>

<1% -

<https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-018-1739-8>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8066912/>

<1% - <https://pubmed.ncbi.nlm.nih.gov/34193055/>

<1% -

https://www.researchgate.net/publication/371180787_Adherence_and_barriers_of_prophylactic_iron_and_folic_acid_supplementation_in_children_a_community-based_mixed_method_study

<1% - https://link.springer.com/chapter/10.1007/978-1-4615-0153-4_11

<1% - <https://www.sciencedirect.com/science/article/pii/S1309104216302719>

<1% - <https://www.nature.com/articles/s41598-023-42603-2>

<1% - <https://pubmed.ncbi.nlm.nih.gov/37482061/>

<1% - <https://pubmed.ncbi.nlm.nih.gov/31205878/>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7335819/>

<1% -

https://www.researchgate.net/publication/357351950_Maternal_Factors_Affecting_the_Incidence_of_Low_Birth_Weight_LBW_in_Indonesia

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10420931/>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9978095/>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5425583/>

<1% - <https://pubmed.ncbi.nlm.nih.gov/34772744/>

<1% - <https://www.sciencedirect.com/science/article/pii/S0306453022002372>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5701713/>

<1% - <https://worldnutrijournal.org/OJS/index.php/WNJ/article/view/260>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5890059/>

<1% -

<https://medicinetoday.com.au/mt/2019/september/feature-article/iron-deficiency-anaemia-children-practical-guide-management>

<1% - <https://pubmed.ncbi.nlm.nih.gov/35745272/>

<1% - <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0271509>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4193389/>

<1% - <https://journal.unnes.ac.id/nju/index.php/kemas/article/view/40875>

<1% - <https://bmcnutr.biomedcentral.com/articles/10.1186/s40795-022-00578-9>

<1% -

<https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-12710-2>

<1% -

<https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-022-14881-4>

<1% - <https://hrr.w.uib.no/files/2019/01/Uddin-2016.pdf>

<1% - <https://psycnet.apa.org/record/2014-24028-008>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6852555/>

<1% -

<https://www.editage.com/insights/ethical-declarations-that-authors-should-provide-at-the-journal-submission-stage>

<1% - <https://mba.nus.edu.sg/admissions-and-funding/>

<1% - <https://pubmed.ncbi.nlm.nih.gov/34955074/>